Mandeville Pharmacy 1655 Highway 190 Mandeville, LA 70448-3582 Phone: (985) 778-2306

Fax: (985) 778-2404

Sep 5 2020 10:34AM Receipt # 39457 Cashier: NORMAN S Drawer #: POS01 - 3342

Rx 44994-00 - JEFFREY BODIN

Items:1

72.99 F

Subtotal 72.99 3.57 Tax Total Total 76.56

Visa Tendered Acct #...1006

> Approval #...033411 Trans ID#...488670

76.56

F = FLEX/FSA/HRA Eligible FSA Eligible \$76.56

Signature required for the following:

- Receipt of Prescriptions

- Receipt of Privacy Practices

Improcied By COVID-19

Signature Captured Electronically

I agree to pay above total amount according to card issuer agreement.

MID:

9139901 1790344

Term ID: App:

US DEBIT

AID:

A0000000980840

AC:

ARQC B80E9D249AD25042

Entry:

INSERT

CVM:

Pin Verified

DebitSale



Scan This QR Code for Patient Specific Medication Information

"Where the Customer is King/Queen"



N0044994-00-00
JEFFREY BODIN
RXLOC
Price 72.99
09/05/2020 Sig:
Sia.

NS

N0044994 09/05/2020 JEFFREY BODIN NORMAN SMITH DEXTROAMP-AMPHETAMIN 10 MG TA

	Mandeville 1655 Hwy	190 • Mandeville, LA 70448	
	\longrightarrow Pharmacy 98!	5-778-2306 	
	JEFFREY BODIN		
	528 BEAU CHENE DR (985) 272-8989		
	MANDEVILLE, LA 70471-1777 PP: RXLO		
	Rx: N0044994 00 09/05/2020	\$72.99 Rph:NS	
8	DEXTROAMP-AMPHETAMIN 10 MG TAB 0055	55-0972-02 Qtv 90	

(985) 778-2306

(985) 272-8989



985-778-2306

Qty 90 PP: RXLOC

Rx: N0044994 00 JEFFREY BODIN 528 BEAU CHENE DR

MANDEVILLE, LA 70471-1777

09/05/2020

(985) 272-8989



DOB: 05/22/1997

Dr. Kevin Mclaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:

NO REFILLS

Rph: NORMAN SMITH (985) 867-5494

00555-0972-02 PHCY ID: 1548616998 Copay

\$72.99

MANDEVILLE PHARMACY

1655 HIGHWAY 190 MANDEVILLE, LA 70448-3582

Rx: N0044994 00 JEFFREY BODIN 528 BEAU CHENE DR

MANDEVILLE, LA 70471-1777 Dr. Kevin Mclaughlin DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth: NO REFILLS

Rph: NORMAN SMITH (985) 867-5494 00555-0972-02 Qty 90 PHCY ID: 1548616998

DUPLICATE RECEIPT \$72.99 Copay

Qty 90

09/05/2020

PP: RXLOC

Call your Doctor for medical advice about side effects. You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"

PRESCRIPTION NOTES

MedGuide™ information for your health®

Mandeville Pharmacy 1655 Highway 190 Mandeville, LA 70448-3582 For: JEFFREY BODIN

Rx: 44994

Date: September 05, 2020 Message ID: 38868793-407

MEDICATION GUIDE

Adderall® (ADD-ur-all) CII

(Dextroamphetamine Saccharate, Amphetamine Aspartate, Dextroamphetamine Sulfate and Amphetamine Sulfate Tablets)

Rx only

Read the Medication Guide that comes with Adderall® before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with Adderall®.

What is the most important information I should know about Adderall®?

The following have been reported with use of Adderall® and other stimulant medicines.

1. Heart-Related problems:

- · sudden death in patients who have heart problems or heart defects
- · stroke and heart attack in adults
- increased blood pressure and heart rate

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting Adderall®.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with Adderall®.

Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Adderall®.

2. Mental (Psychiatric) problems:

All Patients

- new or worse behavior and thought problems
- · new or worse bipolar illness
- new or worse aggressive behavior or hostility

Children and Teenagers

 new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking Adderall®, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

- **3. Circulation problems in fingers and toes** [Peripheral vasculopathy, including Raynaud's phenomenon]:
- fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes.

Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking Adderall®.

What is Adderall®?

Adderall® is a central nervous system stimulant prescription medicine. It is used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD). Adderall® may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD. Adderall® should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Adderall® is also used in the treatment of a sleep disorder called narcolepsy.

Adderall® is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep Adderall® in a safe place to prevent misuse and abuse. Selling or giving away Adderall® may harm others, and is against the law.

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take Adderall®?

Adderall® should not be taken if you or your child:

- · have heart disease or hardening of the arteries
- have moderate to severe high blood pressure
- · have hyperthyroidism
- have an eye problem called glaucoma
- are very anxious, tense, or agitated
- · have a history of drug abuse
- are taking or have taken within the past 14 days an anti-depression medicine called a monoamine

oxidase inhibitor or MAOI.

• are sensitive to, allergic to, or had a reaction to other stimulant medicines Adderall® is not recommended for use in children less than 3 years old.

Adderall® may not be right for you or your child. Before starting Adderall® tell your or your child's doctor about all health conditions (or a family history of) including:

- · heart problems, heart defects, high blood pressure
- · mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- liver or kidney problems

- · thyroid problems
- · seizures or have had an abnormal brain wave test (EEG)
- · circulation problems in fingers or toes

Tell your doctor if you or your child are pregnant, planning to become pregnant, or breastfeeding.

Can Adderall® be taken with other medicines?

Tell your doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements. Adderall® and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking Adderall®.

Your doctor will decide whether Adderall® can be taken with other medicines.

Especially tell your doctor if you or your child take:

- anti-depression medicines including MAOIs
- blood pressure medicines
- · seizure medicines
- · blood thinner medicines
- · cold or allergy medicines that contain decongestants
- · stomach acid medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking Adderall® without talking to your doctor first.

How should Adderall® be taken?

- Take Adderall® exactly as prescribed. Your doctor may adjust the dose until it is right for you or your child.
- Adderall[®] tablets are usually taken two to three times a day. The first dose
 is usually taken when you first wake in the morning. One or two more doses
 may be taken during the day, 4 to 6 hours apart.
- Adderall[®] can be taken with or without food.
- From time to time, your doctor may stop Adderall® treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure
 while taking Adderall[®]. Children should have their height and weight checked
 often while taking Adderall[®]. Adderall[®] treatment may be stopped if a
 problem is found during these check-ups.
- If you or your child take too much Adderall® or overdoses, call your doctor or poison control center right away, or get emergency treatment.

What are possible side effects of Adderall®?

See "What is the most important information I should know about Adderall®?" for information on reported heart and mental problems.

Other serious side effects include:

- · slowing of growth (height and weight) in children
- · seizures, mainly in patients with a history of seizures
- · eyesight changes or blurred vision

Common side effects include:

- · stomach ache
- decreased appetite
- nervousness

Adderall® may affect your or your child's ability to drive or do other dangerous activities.

Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist

for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Adderall®?

- Store Adderall[®] in a safe place at room temperature, 20° to 25°C (68° to 77°F)
- Keep Adderall® and all medicines out of the reach of children.

General information about Adderall®

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Adderall® for a condition for which it was not prescribed. Do not give Adderall® to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about Adderall®. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Adderall® that was written for healthcare professionals. For more information about Adderall®, please contact Teva Pharmaceuticals at 1-888-838-2872.

What are the ingredients in Adderall®?

Active Ingredient: dextroamphetamine saccharate, amphetamine aspartate monohydrate, dextroamphetamine sulfate, USP, amphetamine sulfate, USP

Inactive Ingredients: lactitol, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, and other ingredients.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Teva Select Brands, Horsham, PA 19044

Division of Teva Pharmaceuticals USA

Revised October 2015