


YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	A. WEISS, DO	DRUG DESCRIPTION	
BIRTH DATE	05/22/97				
MEDICATION	D-AMPHETAMINE SALT COMBO 30MG TABS				
QUANTITY	90	PATIENT ALLERGIES	<ALLERGY HISTORY NOT KNOWN>		
DIRECTIONS	TAKE 1 TABLET BY MOUTH THREE TIMES DAILY				PEACH
					FRONT: b / 974
					BACK: 30

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take longer than you have been told by your doctor. Use this drug only as you were told by your doctor. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death. If you have any questions, talk with your doctor.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: if you have an allergy to dextroamphetamine, amphetamine, or any other part of this drug. **TELL YOUR DOCTOR:** if you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. **TELL YOUR DOCTOR:** if you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** if you have any of these health problems: Glaucoma; nervous, anxious, or tense state; or overactive thyroid. **TELL YOUR DOCTOR:** if you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** if you have kidney disease. **TELL YOUR DOCTOR:** if you are taking acetazolamide. **TELL YOUR DOCTOR:** if you are taking sodium bicarbonate. **TELL YOUR DOCTOR:** if you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. **TELL YOUR DOCTOR:** if you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** if you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC), natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow directions and instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. To gain the most benefit, do not miss doses. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, decongestants or like products, and some natural products or aids. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down the toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. This drug may be habit-forming with long-term use. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may have some heart tests before starting this drug. Talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Talk with the doctor. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Do not switch between this drug and other products that have the same drugs in them without checking with the doctor. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like low mood (depression) or bipolar illness, or if a family member has killed themselves. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like low mood (depression), thoughts of killing yourself, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A very bad and sometimes deadly health problem called serotonin syndrome may happen if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea, upset stomach, or throwing up; or very bad headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. A fast heartbeat. Very nervous and excitable. Not able to get or keep an erection. Change in sex interest. A heartbeat that does not feel normal. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis)

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188 12T

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777
(985)272-8989

RX # 2170156-05382

DATE: 05/26/19

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS

New NDC: 00555-0974-02

\$ 0.00

A. WEISS, DO
MFG: TEVA
AAT/MRA/MRA/SVV/CKM/TCH/ELL

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 191464627522112999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777
(985)272-8989

RX # 2170156-05382

DATE: 05/26/19

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS

New NDC: 00555-0974-02

\$ 0.00

A. WEISS, DO
MFG: TEVA
AAT/MRA/MRA/SVV/CKM/TCH/ELL

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 191464627522112999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

WAITING

SUN 1:05PM

New


D-AMPHETAMINE SALT COMBO 30MG TABS

00555-0974-02

SAFE

QTY 90

20 DRAM



PEACH

FRONT: b / 974

BACK: 30

AAT/MRA/MRA/SVV/CKM/TCH/ELL

Med Guide