



705WB900721

P.O. Box 29063, Phoenix, AZ 85038-9063 (Store 21147)
NABP #: 0323624

Received

4/18/19

www.walgreens.com

It is easy and quick to pay using your credit card either online at www.walgreens.com or by calling our automated telephone system 1-877-250-5823. Those who are deaf or hard of hearing can call TTY 1-800-573-1833.

Ship to: JEFFREY BODIN 528 BEAU CHENE DRIVE MANDEVILLE, LA 70471	Account Balance: \$0.00 Invoice Number: WB9DD721 Shipment Date: 4/12/2019 Member ID: *****2043
--	---

Received 4/18/19

Start of Prescription on 4/20/19

Prescription Items

Rx Number	Item Description	Quantity	Amount
1785907-05823	BUPROPION XL 300MG TABLETS ACTAVIS 45963-0142-05	30	0.00

THANK YOU FOR YOUR PROMPT PAYMENT.
All credit card charges are pending authorization.

Current Order:	0.00
Previous Balance:	0.00
Amount Due:	0.00

- Thank you for your order. Please visit us again at www.walgreens.com for all your pharmacy needs.
- Please make us aware if you have any updates regarding your drug allergies or health conditions by calling our customer service number or updating your profile online



Personal Prescription Information

Patient: JEFFREY BODIN

RX#: 1785907 - 05823

Drug: BUPROPION XL 300MG TABLETS

Dr: WEISS

Directions: TAKE 1 TABLET BY MOUTH EVERY DAY IN THE MORNING

IF YOU HAVE ANY QUESTIONS ABOUT YOUR MEDICATION AFTER YOU READ THIS INFORMATION, WE URGE YOU TO CONTACT OUR PHARMACISTS AT www.walgreens.com OR BY CALLING TOLL-FREE 1-877-250-5823.

The following is the generic name for your medication; however, this does not imply that a generic substitution is available.

GENERIC NAME: Bupropion Extended-Release Tablets (byoo PROE pee on)

WARNING: Drugs like this one have raised the chance of suicidal thoughts or actions in children and young adults. The risk may be greater in people who have had these thoughts or actions in the past. All people who take this drug need to be watched closely. Call the doctor right away if signs like low mood (depression), nervousness, restlessness, grouchiness, panic attacks, or changes in mood or actions are new or worse. Call the doctor right away if any thoughts or actions of suicide occur. **COMMON USES:** It is used to treat low mood (depression). It is used to treat seasonal affective disorder (SAD). It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to bupropion or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; or shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. **TELL YOUR DOCTOR:** If you have ever had seizures. **TELL YOUR DOCTOR:** If you drink a lot of alcohol and you stop drinking all of a sudden. **TELL YOUR DOCTOR:** If you use certain other drugs like drugs for seizures or anxiety and you stop using them all of a sudden. **TELL YOUR DOCTOR:** If you have ever had an eating problem like anorexia or bulimia. **TELL YOUR DOCTOR:** If you have any of these health problems: Kidney disease or liver disease. **TELL YOUR DOCTOR:** If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are taking another drug that has the same drug in it. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Do not take this drug more often than you are told. This may raise the risk of seizures. Be sure you know how far apart to take your doses. Take in the morning if taking once a day. Take with or without food. If you are not able to sleep, do not take this drug too close to bedtime. Talk with your doctor. Swallow whole. Do not chew, break, or crush. To gain the most benefit, do not miss doses. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. If you have trouble swallowing, talk with your doctor. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain

unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: For all patients taking this drug: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert or have clear eyesight until you see how this drug affects you. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not stop taking this drug all of a sudden without calling your doctor. You may have a greater risk of side effects. If you need to stop this drug, you will want to slowly stop it as ordered by your doctor. High blood pressure has happened with this drug. Have your blood pressure checked as you have been told by your doctor. This drug may raise the chance of seizures. The chance may be higher in people who have certain health problems, use certain other drugs, or drink a lot of alcohol. Talk to your doctor to see if you have a greater chance of seizures while taking this drug. Avoid drinking alcohol while taking this drug. Talk with your doctor before you use other drugs and natural products that slow your actions. It may take several weeks to see the full effects. Some people may have a higher chance of eye problems with this drug. Your doctor may want you to have an eye exam to see if you have a higher chance of these eye problems. Call your doctor right away if you have eye pain, change in eyesight, or swelling or redness in or around the eye. If you are taking digoxin, talk with your doctor. You may need to have your blood work checked more closely while you are taking it with this drug. This drug is not approved for use in children. Talk with the doctor. If you are 65 or older, use this drug with care. You could have more side effects. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby. If you smoke: Not all products are approved for use to help stop smoking. Talk with the doctor to make sure that you have the right product. New or worse mental, mood, or behavior problems have happened when bupropion has been used to stop smoking. These problems include thoughts of suicide or killing someone else, depression, forceful actions, fury, anxiety, and anger. These problems have happened in people with and without a history of mental or mood problems. Talk with the doctor.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Change in how you act. Feeling confused. Hallucinations (seeing or hearing things that are not there). If seizures are new or worse after starting this drug. A big weight gain or loss. Chest pain or pressure or a fast heartbeat. A heartbeat that does not feel normal. Swelling. Shortness of breath. Change in hearing. Ringing in ears. Passing urine more often. Swollen gland. Trouble moving around. Very bad muscle or joint pain. A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Dizziness. Headache. Belly pain. Shakiness. Feeling nervous and excitable. Strange or odd dreams. Upset stomach or throwing up. Constipation. Gas. Dry mouth. Not able to sleep. Muscle or joint pain. Nose or throat irritation. Sweating a lot. Not hungry. A change in weight without trying. For some brands, you may see the tablet shell in your stool. For these brands, this is normal and not a cause for concern. If you have questions, talk with your doctor. These are not all of the side effects that may occur. If

BUPROPION XL 300MG TABLETS – page 1 of 2

Notice: Your prescription order may be processed or filled by another Walgreens pharmacy, such as Walgreens.com pharmacy or, part of our network of pharmacies under common ownership may process or fill the order as well. Written information about your prescription has been provided for you. Please read this information before you take the medication. If you have questions concerning this prescription, a pharmacist is available during normal business hours to answer these questions at 1-800-Walgreens.

BUPROPION XL 300MG TABLETS

you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

HOW TO GET RID OF UNUSED MEDICINE:

Do not flush unused medications or pour down a sink or drain. For detailed instructions on how to properly get rid of unwanted medicine, visit the Food and Drug Administration's (FDA's) website at: www.fda.gov/forconsumers/consumerupdates/ucm101653.htm

MEDICATION GUIDE

BUPROPION SR (ByoU-PROF-pee-on)

(bupropion hydrochloride)
Sustained-Release Tablets

IMPORTANT: Be sure to read the three sections of this Medication Guide. The first section is about the risk of suicidal thoughts and actions with antidepressant medicines; the second section is about the risk of changes in thinking and behavior, depression and suicidal thoughts or actions with medicines used to quit smoking; and the third section is about Bupropion SR.[®]

Antidepressant Medicines, Depression and Other Serious

Mental Illnesses, and Suicidal Thoughts or Actions

This section of the Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines.

What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions?

1. Antidepressant medicines may increase the risk of suicidal thoughts or actions in some children, teenagers, or young adults within the first few months of treatment.

2. Depression or other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions. These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.

3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?

• Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.

• Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.

• Keep all follow-up visits with your healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call your healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:

• thoughts about suicide or dying

• attempts to commit suicide

• new or worse depression

• new or worse anxiety

• feeling very agitated or restless

• panic attacks

• trouble sleeping (insomnia)

• new or worse irritability

• acting aggressive, being angry, or violent

• acting on dangerous impulses

• an extreme increase in activity and talking (mania)

• other unusual changes in behavior or mood

What else do I need to know about antidepressant medicines?

• **Never stop an antidepressant medicine without first talking to a healthcare provider.** Stopping an antidepressant medicine suddenly can cause other symptoms.

• **Antidepressants are medicines used to treat depression and other illnesses.** It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.

• **Antidepressant medicines have other side effects.** Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.

• **Antidepressant medicines can interact with other medicines.** Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with your healthcare provider.

It is not known if bupropion SR is safe and effective in children under the age of 18.

Quitting Smoking, Quit-Smoking Medications, Changes in Thinking and Behavior, Depression, and Suicidal Thoughts or Actions

This section of the Medication Guide is only about the risk of changes in thinking and behavior, depression and suicidal thoughts or actions with drugs used to quit smoking. Although bupropion SR is not a treatment for quitting smoking, it contains the same active ingredient (bupropion hydrochloride) as ZYBAN[®] which is used to help patients quit smoking.

Talk to your healthcare provider or your family member's healthcare provider about:

• all risks and benefits of quit-smoking medicines.

• all treatment choices for quitting smoking.

When you try to quit smoking, with or without bupropion, you may have symptoms that may be due to nicotine withdrawal, including:

• urge to smoke

• frustration

• depressed mood

• trouble sleeping

• irritability

• restlessness

• weight gain

Some people have even experienced suicidal thoughts when trying to quit smoking without medication. Sometimes quitting smoking can lead to worsening of mental health problems that you already have, such as depression.

Some people have had serious side effects while taking bupropion to help them quit smoking, including: **New or worse mental health problems, such as changes in behavior or thinking, aggression, hostility, agitation, depression, or suicidal thoughts or actions.** Some people had these symptoms when they began taking bupropion, and others developed them after several weeks of treatment, or after stopping bupropion. These symptoms

happened more often in people who had a history of mental health problems before taking bupropion than in people without a history of mental health problems.

Stop taking bupropion SR, and call your healthcare provider right away if you, your family, or caregiver notice any of these symptoms. Work with your healthcare provider to decide whether you should continue to take bupropion SR. In many people, these symptoms went away after stopping bupropion SR, but in some people symptoms continued after stopping bupropion SR. It is important for you to follow-up with your healthcare provider until your symptoms go away.

Before taking bupropion SR, tell your healthcare provider if you have ever had depression or other mental health problems. You should also tell your healthcare provider about any symptoms you had during other times you tried to quit smoking, with or without bupropion.

What Other Important Information Should I Know About Bupropion SR?

• **Seizures: There is a chance of having a seizure (convulsion, fit) with bupropion SR, especially in people:**

o with certain medical problems.

o who take certain medicines.

The chance of having seizures increases with higher doses of bupropion SR. For more information, see the sections "Who should not take bupropion SR?" and "What should I tell my healthcare provider before taking bupropion SR?." Tell your healthcare provider about all of your medical conditions and all the medicines you take. Do not take any other medicines while you are taking bupropion SR unless your healthcare provider has said it is okay to take them.

If you have a seizure while taking bupropion SR, stop taking the tablets and call your healthcare provider right away. Do not take bupropion SR again if you have a seizure.

• **High blood pressure (hypertension). Some people get high blood pressure, that can be severe, while taking bupropion SR.**

The chance of high blood pressure may be higher if you also use nicotine replacement therapy (such as a nicotine patch) to help you stop smoking (see the section of this Medication Guide called "How should I take bupropion SR?").

• **Manic episodes.** Some people may have periods of mania while taking bupropion SR, including:

o Greatly increased energy

o Severe trouble sleeping

o Racing thoughts

o Reckless behavior

o Unusually grand ideas

o Excessive happiness or irritability

o Talking more or faster than usual

If you have any of the above symptoms of mania, call your healthcare provider.

• **Unusual thoughts or behaviors.** Some patients have unusual thoughts or behaviors while taking bupropion SR, including delusions (believe you are someone else), hallucinations (seeing or hearing things that are not there), paranoia (feeling that people are against you), or feeling confused. If this happens to you, call your healthcare provider.

• Visual problems.

o eye pain

o swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

• **Severe allergic reactions. Some people can have severe allergic reactions to bupropion SR. Stop taking bupropion SR and call your healthcare provider right away if you get a rash, itching, hives, fever, swollen lymph glands, painful sores in the mouth or around the eyes, swelling of the lips or tongue, chest pain, or have trouble breathing.** These could be signs of a serious allergic reaction.

What is bupropion SR?

Bupropion SR is a prescription medicine used to treat adults with a certain type of depression called major depressive disorder. **Who should not take bupropion SR if you**

• have or had a seizure disorder or epilepsy.

• have or had an eating disorder such as anorexia nervosa or bulimia.

• **are taking any other medicines that contain bupropion, including ZYBAN (used to help people stop smoking), WELLBUTRIN[®], WELLBUTRIN XL[®], APLENZIN[®], or FORFIVO XL[®].** Bupropion is the same active ingredient that is in WELLBUTRIN SR.

• drink a lot of alcohol and abruptly stop drinking, or use medicines called sedatives (these make you sleepy), benzodiazepines, or anti-seizure medicines, and you stop using them all of a sudden.

• take a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.

o do not take an MAOI within 2 weeks of stopping bupropion SR unless directed to do so by your healthcare provider.

o do not start bupropion SR if you stopped taking an MAOI in the last 2 weeks unless directed to do so by your healthcare provider.

• are allergic to the active ingredient in bupropion SR, bupropion, or to any of the inactive ingredients. If you need more information about the ingredients in this medicine ask your pharmacist for a complete list.

What should I tell my healthcare provider before taking bupropion SR?

Tell your healthcare provider if you have ever had depression, suicidal thoughts or actions, or other mental health problems. See "Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal Thoughts or Actions."

• **Tell your healthcare provider about your other medical conditions including if you:**

o have liver problems, especially cirrhosis of the liver.

o have kidney problems.

o have, or have had, an eating disorder, such as anorexia nervosa or bulimia.

o have had a head injury.

o have had a seizure (convulsion, fit).

o have a tumor in your nervous system (brain or spine).

o are a diabetic taking insulin or other medicines to control your blood sugar.

o drink alcohol.

o abuse prescription medicines or street drugs.

o are pregnant or plan to become pregnant.

o are breastfeeding. Bupropion SR passes into your milk in small amounts.

Tell your healthcare provider about all the medicines you take, including prescription, over-the-counter medicines, vitamins, and herbal supplements. Many medicines increase your chances of having seizures or other serious side effects if you take them while you are taking bupropion SR.

How should I take bupropion SR?

• Take bupropion SR exactly as prescribed by your healthcare provider. Do not change your dose or stop taking bupropion SR without talking with your healthcare provider first.

• **Swallow bupropion SR tablets whole. Do not chew, cut, or crush bupropion SR tablets.** If you do, the medicine will be more likely to get side effects including seizures. **Tell your healthcare provider if you cannot swallow tablets.**

• Bupropion SR tablets may have an odor. This is normal.

• Take bupropion SR at the same time each day.

• Take your doses of bupropion SR at least 8 hours apart.

• If you miss a dose, do not take an extra dose to make up for the dose you missed. Wait and take your next dose at the regular time. **This is very important.** Too much bupropion SR can increase your chance of having a seizure.

• If you take too much bupropion SR, or overdose, call your local emergency room or poison control center right away.

• **Do not take any other medicines while taking bupropion SR unless your healthcare provider has told you it is okay.**

• If you are taking bupropion SR for the treatment of major depressive disorder, it may take several weeks for you to feel that bupropion SR is working. Once you feel better, it is important to keep taking bupropion SR exactly as directed by your healthcare provider. Call your healthcare provider if you do not feel bupropion SR is working for you.

What should I avoid while taking bupropion SR?

• Limit or avoid using alcohol during treatment with bupropion SR. If you usually drink a lot of alcohol, talk with your healthcare provider before suddenly stopping. If you suddenly stop drinking alcohol, you may increase your risk of having seizures.

• Do not drive a car or use heavy machinery until you know how bupropion SR affects you. Bupropion SR can affect your ability to do these things safely.

What are possible side effects of bupropion SR?

Bupropion SR can cause serious side effects. See the sections at the beginning of this Medication Guide for information about serious side effects of bupropion SR.

The most common side effects of bupropion SR include:

• headache

• dizziness

• constipation

• dry mouth

• trouble sleeping

• sore throat

• nausea

If you have nausea, take your medicine with food. If you have trouble sleeping, do not take your medicine too close to bedtime. Tell your healthcare provider right away about any side effects that bother you.

These are not all the possible side effects of bupropion SR. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store bupropion SR sustained-release tablets?

• Store bupropion SR sustained-release tablets at room temperature between 59°F and 86°F (15°C to 30°C).

• Keep bupropion SR dry and out of the light.

Keep bupropion SR and all medicines out of the reach of children.

General information about the safe and effective use of bupropion SR.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use bupropion SR for a condition for which it was not prescribed. Do not give bupropion SR to other people, even if they have the same symptoms you have. It may harm them.

If you take a urine drug screening test, bupropion SR may make the test result positive for amphetamines. If you tell the person giving you the drug screening test that you are taking bupropion SR, they can do a more specific drug screening test that should not have this problem.

This Medication Guide summarizes important information about bupropion SR. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about bupropion SR that is written for healthcare professionals.

What are the ingredients in bupropion SR sustained-release tablets?

Active ingredient: bupropion hydrochloride.

Inactive ingredients in this medicine ask your pharmacist for a complete list.

This Medication Guide has been approved by the U.S. Food and Drug Administration. Revised 05/2017