

Attn: Ashley L Weiss DO + Grinasha M Dillon // JEFFREY THOMAS BODIN // 2020-11-10

Attn: Ashley L Weiss DO + Grinasha M Dillon // JEFFREY THOMAS BODIN // 2020-11-10

Fax TO,
1-504-988-0302

Attached,
2020-11-10 Dr Weiss - fax - Adderall 40 mg prescription updates_ENTER INTO MEDICAL
SYSTEM (1)

(1)
Social Security Administration,
SSI Statement of Benefits

(2)
Jul 2020 -> Nov 2020,
Adderall 40 mg records

Jeffrey Bodin

2020-11-10 Dr Weiss - fax - Adderall 40 mg prescription updates_ENTER INTO MEDICAL SYSTEM

Jeffrey Thomas Bodin
528 Beau Chene Dr
Mandeville, LA 70471

Grinasha M Dillon
4000 Bienville St
Ste G, New Orleans, LA 70119
P: 504-988-0306

CC:

Dr. Ashley L. Weiss DO
4000 Bienville St
Ste G, New Orleans, LA 70119

aweiss3@tulane.edu

P: 504-988-0301 // F: 504-988-0302

FAX TO: 504-988-0302


Mrs. Grinasha,

1. Enter attached prescription filled records.
 - a. Into Tulane's medical records system.
 - b. Plus, any other system accessible.
 - c. Scanning them in.
2. Scan into records the "SSI Benefit Verification Letter"
 - a. For assisting in solving future collections notices.
 - i. By Tulane / Corporate / Your clinics "billing" in some form
 - ii. Mistakenly sent, I'm sure.
 - iii. As it resolved in Sep/Oct. On mailing my Cease/Desist.
3. Attached: SSI Ltr, Prescription Jul 2020 – Nov 2020 40 mg.



Social Security Administration Benefit Verification Letter

Date: November 10, 2020
BNC#: 20UC561E10803
REF: DI


JEFFREY THOMAS BODIN
528 BEAU CHENE DR
MANDEVILLE LA 70471-1777

0101BEV8S097IUJ CCM.M72.BEV8S.R201111

You asked us for information from your record. The information that you requested is shown below. If you want anyone else to have this information, you may send them this letter.

Information About Supplemental Security Income Payments

Beginning October 2020, the current Supplemental Security Income payment is \$783.00.

This payment amount may change from month to month if income or living situation changes.

Supplemental Security Income Payments are paid the month they are due.

(For example, Supplemental Security Income Payments for March are paid in March.)

We found that you became disabled under our rules on May 22, 2015.

Type of Supplemental Security Income Payment Information

You are entitled to monthly payments as a disabled individual.

Date of Birth Information

The date of birth shown on our records is May 22, 1997.

Suspect Social Security Fraud?

Please visit <http://oig.ssa.gov/r> or call the Inspector General's Fraud Hotline at 1-800-269-0271 (TTY 1-866-501-2101).

See Next Page

If You Have Questions

We invite you to visit our web site at www.socialsecurity.gov on the Internet to find general information about Social Security. If you have any specific questions, you may call us toll-free at 1-800-772-1213, or call your local office at 1-866-887-8997. We can answer most questions over the phone. If you are deaf or hard of hearing, you may call our TTY number, 1-800-325-0778. You can also write or visit any Social Security office. The office that serves your area is located at:

SOCIAL SECURITY
64285 HIGHWAY 434
LACOMBE LA 70445

If you do call or visit an office, please have this letter with you. It will help us answer your questions. Also, if you plan to visit an office, you may call ahead to make an appointment. This will help us serve you more quickly when you arrive at the office.

Social Security Administration

#1500

Rx#: 2224291 07/03/20
T/P Auth 9: 484615007035G
BODIN, JEFFREY
528 BEAUVU CHENE DR
MANDEVILLE, LA 70471
MCLAUGHLIN, KEVIN EARL MD
AMPHETAMINE 10 MG
For > ADDERALL 10 MG
NDC: 13107-0070-01
Disp Qty: 90.000TAB

DOB: 05/22/97
Ph: (985) 272-8989
DL #: 010591503
Refills: NR by 10/01/20
TAB AUROBINDO



B

YOUR COST: \$33.37

#1500

Rx#: 2224291 07/03/20
Tx: 1713411
Carrier: ENV
DS: 30
NEW

THIS MEDICINE IS A(N) LIGHT BLUE
DOUBLE-SCORED ROUND-SHAPED TABLET
IMPRINTED WITH U27. ADD INFO: MOTTLED
APPEARANCE CONVEX LIGHT BLUE TO BLUE

CASH PRICE: \$120.99
YOUR COST: \$33.37

INFO: 1500 13107007001 OPT: 1500 664 185 0620201

79716

**Please see
the accompanying
pages
for more information...**

To stop receiving sponsored info at our pharmacy, call 888-336-5744 or visit remove-me.net Use code: 15006664 185 0620201

YOUR PERSONAL PRESCRIPTION INFORMATION

Winn-Dixie
+ Pharmacy
#1500

If you have any questions, please contact your Winn-Dixie pharmacist at
PHONE # (985) 893-0187

PATIENT MEDICATION DIRECTIONS:

TAKE ONE TABLET BY MOUTH THREE TIMES A DAY

BODIN, JEFFREY

AMPHETAMINE 10 MG TAB

GENERIC NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death. **COMMON USES:** It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. **TELL YOUR DOCTOR:** If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, ibuprofen or like products, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or

peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable. Headache. Trouble sleeping. Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

This information should not be used to decide whether or not to take this medicine or any other medicine. Only your health care provider has the knowledge and training to decide which medicines are right for you. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from your health care provider. You must talk with your healthcare provider for complete information about the risks and benefits of using this medicine. Copyright 2020 CDI, LLC. All rights reserved. All rights reserved. Database Edition 20.2.1.003 Information Expires July 22, 2020

MEDICATION GUIDE

DEXTROAMPHETAMINE SACCHARATE, AMPHETAMINE ASPARTATE, DEXTROAMPHETAMINE SULFATE AND AMPHETAMINE SULFATE

(DEX-troe-am-FET-uh-meen SACK-uh-rate, am-FET-uh-meen ass-PAR-tate, DEX-troe-am-FET-uh-meen SULL-fate, and am-FET-uh-meen SULL-fate)

TABLETS (MIXED SALTS OF A SINGLE ENTITY AMPHETAMINE PRODUCT) CII

Read the Medication Guide that comes with dextroamphetamine saccharate, amphetamine aspartate, dextroamphetamine sulfate and amphetamine sulfate tablets (mixed amphetamine salts) before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with mixed amphetamine salts.

What is the most important information I should know about mixed amphetamine salts?

The following have been reported with use of mixed amphetamine salts and other stimulant medicines.

1. Heart-related problems:

- sudden death in patients who have heart problems or heart defects
- stroke and heart attack in adults
- increased blood pressure and heart rate

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting mixed amphetamine salts.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with mixed amphetamine salts.

Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking mixed amphetamine salts.

2. Mental (Psychiatric) problems:

All Patients

- new or worse behavior and thought problems
- new or worse bipolar illness
- new or worse aggressive behavior or hostility

Children and Teenagers

- new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking mixed amphetamine salts, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

3. Circulation problems in fingers and toes [Peripheral Vasculopathy, Including Raynaud's Phenomenon]:

- Fingers or toes may feel numb, cool, painful
- Fingers or toes may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in your fingers or toes.

Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking mixed amphetamine salts.

What are mixed amphetamine salts?

Mixed amphetamine salts are a central nervous system stimulant prescription medicine. It is used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD). Mixed amphetamine salts may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD.

Mixed amphetamine salts should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Mixed amphetamine salts are also used in the treatment of a sleep disorder called narcolepsy.

Mixed amphetamine salts are a federally controlled substance (CII) because it can be abused or lead to dependence. Keep mixed amphetamine salts in a safe place to prevent misuse and abuse. Selling or giving away mixed amphetamine salts may harm others, and is against the law.

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take mixed amphetamine salts?

Mixed amphetamine salts should not be taken if you or your child:

- have heart disease or hardening of the arteries
- have moderate to severe high blood pressure
- have hyperthyroidism
- have an eye problem called glaucoma
- are very anxious, tense, or agitated
- have a history of drug abuse
- are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOI.
- are sensitive to, allergic to, or had a reaction to other stimulant medicines

Mixed amphetamine salts are not recommended for use in children less than 3 years old.

Mixed amphetamine salts may not be right for you or your child. Before starting mixed amphetamine salts tell your or your child's doctor about all health conditions (or a family history of) including:

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome

- liver or kidney problems
- circulation problems in fingers and toes
- thyroid problems
- seizures or have had an abnormal brain wave test (EEG)

Tell your doctor if you or your child are pregnant, planning to become pregnant, or breastfeeding.

Can mixed amphetamine salts be taken with other medicines?

Tell your doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements. Mixed amphetamine salts and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking mixed amphetamine salts.

Your doctor will decide whether mixed amphetamine salts can be taken with other medicines.

Especially tell your doctor if you or your child take:

- anti-depression medicines including MAOIs
- blood pressure medicines
- seizure medicines
- blood thinner medicines
- cold or allergy medicines that contain decongestants
- stomach acid medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking mixed amphetamine salts without talking to your doctor first.

How should mixed amphetamine salts be taken?

- **Take mixed amphetamine salts exactly as prescribed.** Your doctor may adjust the dose until it is right for you or your child.
- Mixed amphetamine salt tablets are usually taken two to three times a day. The first dose is usually taken when you first wake in the morning. One or two more doses may be taken during the day, 4 to 6 hours apart.
- Mixed amphetamine salts can be taken with or without food.
- From time to time, your doctor may stop mixed amphetamine salts treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure while taking mixed amphetamine salts. Children should have their height and weight checked often while taking mixed amphetamine salts. Mixed amphetamine salts treatment may be stopped if a problem is found during these check-ups.
- **If you or your child take too much mixed amphetamine salts or overdose, call your doctor or poison control center right away, or get emergency treatment.**

What are possible side effects of mixed amphetamine salts?

See “**What is the most important information I should know about mixed amphetamine salts?**” for information on reported heart and mental problems.

Other serious side effects include:

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision
- serotonin syndrome. A potentially life-threatening problem called serotonin syndrome can happen when medicines such

as mixed amphetamine salts are taken with certain other medicines. Symptoms of serotonin syndrome may include:

- agitation, hallucinations, coma or other changes in mental status
- problems controlling your movements or muscle twitching
- fast heartbeat
- high or low blood pressure
- sweating or fever
- nausea or vomiting
- diarrhea
- muscle stiffness or tightness

Common side effects include:

- stomach ache
- decreased appetite
- nervousness

Mixed amphetamine salts may affect your or your child’s ability to drive or do other dangerous activities.

Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store mixed amphetamine salts?

- Store mixed amphetamine salts in a safe place at room temperature, 20° to 25°C (68° to 77°F).
- **Keep mixed amphetamine salts and all medicines out of the reach of children.**

General information about mixed amphetamine salts

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use mixed amphetamine salts for a condition for which it was not prescribed. Do not give mixed amphetamine salts to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about mixed amphetamine salts. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about mixed amphetamine salts that was written for healthcare professionals.

What are the ingredients in mixed amphetamine salts?

Active Ingredient: dextroamphetamine saccharate, amphetamine aspartate, dextroamphetamine sulfate and amphetamine sulfate.

Inactive Ingredients: If you need more information about the inactive ingredients in this medicine ask your pharmacist for a complete list.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Rev. 3/2020

ATTENTION: This is information about the product featured in the leaflet, NOT your prescription medication.

Notice of Privacy Practices

Your Information. Your Rights. Our Responsibilities.

Effective Date: August 7, 2017

This notice describes how medical information about you may be used and disclosed and how you can get access to this information. **Please review it carefully.**

You have the right to:

- Get a copy of your paper or electronic medical record
- Correct your paper or electronic medical record
- Request confidential communication
- Ask us to limit the information we share
- Get a list of those with whom we've shared your information
- Get a copy of this privacy notice
- Choose someone to act for you
- File a complaint if you believe your privacy rights have been violated

Our Uses and Disclosures

We may use and share your information as we:

- Treat you
- Run our organization
- Bill for your services
- Help with public health and safety issues
- Do research
- Comply with the law

Your Choices

You have some choices in the way that we use and share information as we:

- Tell family and friends about your condition
 - Provide disaster relief
 - Include you in a hospital directory
 - Provide mental health care
 - Market our services and sell your information
 - Raise funds
-
- Respond to organ and tissue donation requests
 - Work with a medical examiner or funeral director
 - Address workers' compensation, law enforcement, and other government requests
 - Respond to lawsuits and legal actions

YOUR RIGHTS

When it comes to your health information, you have certain rights. This section explains your rights and some of our responsibilities to help you.

Get an electronic or paper copy of your medical record

- You can ask to see or get an electronic or paper copy of your medical record and other health information we have about you. Ask us how.
- We will provide a copy or a summary of your health information, usually within 30 days of your request. We may charge a reasonable, cost-based fee.

Ask us to correct your medical record

- You can ask us to correct health information about you that you think is incorrect or incomplete. Ask us how to do this.
- We may say "no" to your request, but we'll tell you why in writing within 60 days.

Request confidential communications

- You can ask us to contact you in a specific way (for example, home or office phone) or to send mail to a different address.
- We will say "yes" to all reasonable requests.

Ask us to limit what we use or share

- You can ask us not to use or share certain health information for treatment, payment, or our operations. We are not required to agree to your request, and we may say "no" if it would affect your care.
 - If you pay for a service or health care item out-of-pocket in full, you can ask us not to share that information for the purpose of payment or our operations with your health insurer.
- We will say "yes" unless a law requires us to share that information.

Get a list of those with whom we've shared information

- You can ask for a list (accounting) of the times we've shared your health information for six years prior to the date you ask, who we shared it with, and why.
- We will include all the disclosures except for those about treatment, payment, and health care operations, and certain other disclosures (such as any you asked us to make). We'll provide one accounting a year for free but will charge a reasonable, cost-based fee if you ask for another one within 12 months.

Get a copy of this privacy notice

You can ask for a paper copy of this notice at any time, even if you have agreed to receive the notice electronically. We will provide you with a paper copy promptly.

Choose someone to act for you

- If you have given someone medical power of attorney or if someone is your legal guardian, that person can exercise your rights and make choices about your health information.
- We will make sure the person has this authority and can act for you before we take any action.

File a complaint if you feel your rights are violated

- You can complain if you feel we have violated your rights by contacting the Health Privacy Office, 8928 Prominence Pkwy, Building 200, Jacksonville, Florida 32256 and submitting a written complaint
- You can file a complaint with the U.S. Department of Health and Human Services Office for Civil Rights by sending a letter to 200 Independence Avenue, S.W., Washington, D.C. 20201, calling 1-877-696-6775, or visiting www.hhs.gov/ocr/privacy/hipaa/complaints/.
- We will not retaliate against you for filing a complaint.

YOUR CHOICES

For certain health information, you can tell us your choices about what we share. If you have a clear preference for how we share your information in the situations described below, talk to us. Tell us what you want us to do, and we will follow your instructions.

In these cases, you have both the right and choice to tell us to:

- Share information with your family, close friends, or others involved in your care
- Share information in a disaster relief situation

If you are not able to tell us your preference, for example if you are unconscious, we may go ahead and share your information if we believe it is in your best interest. We may also share your information when needed to lessen a serious and imminent threat to health or safety.

In these cases we never share your information unless you give us written permission.

- Marketing purposes
- Sale of your information
- Most sharing of psychotherapy notes

In the case of fundraising:

- We may contact you for fundraising efforts, but you can tell us not to contact you again.

OUR USES AND DISCLOSURES

How do we typically use or share your health information?

We typically use or share your health information in the following ways:

Treat you: We can use your health information and share it with other professionals who are treating you. *Example: A doctor treating you for an injury asks another doctor about your overall health condition.*

Run our organization: We can use and share your health information to run our practice, improve your care, and contact you when necessary. *Example: We use health information about you to manage your treatment and services.*

Bill for your services: We can use and share your health information to bill and get payment from health plans or other entities. *Example: We give information about you to your health insurance plan so it will pay for your services.*

How else can we use or share your health information?

We are allowed or required to share your information in other ways – usually in ways that contribute to the public good, such as public health and research. We have to meet many conditions in the law before we can share your information for these purposes. For more information see: www.hhs.gov/ocr/privacy/hipaa/understanding/consumers/index.html.

Help with public health and safety issues

We can share health information about you for certain situations such as:

- Preventing disease
- Helping with product recalls
- Reporting adverse reactions to medications
- Reporting suspected abuse, neglect, or domestic violence
- Preventing or reducing a serious threat to anyone's health or safety

Do research

We can use or share your information for health research.

Comply with the law

We will share information about you if state or federal laws require it, including with the Department of Health and Human Services if it wants to see that we're complying with federal privacy law.

Respond to organ and tissue donation requests

We can share health information about you with organ procurement organizations.

Work with a medical examiner or funeral director

We can share health information with a coroner, medical examiner, or funeral director when an individual dies.

Address workers' compensation, law enforcement, and other government requests

We can use or share health information about you:

- For workers' compensation claims
- For law enforcement purposes or with a law enforcement official
- With health oversight agencies for activities authorized by law
- For special government functions such as military, national security, and presidential protective services

Respond to lawsuits and legal actions

We can share health information about you in response to a court or administrative order, or in response to a subpoena.

OUR RESPONSIBILITIES

- We are required by law to maintain the privacy and security of your protected health information.
- We will let you know promptly if a breach occurs that may have compromised the privacy or security of your information.
- We must follow the duties and privacy practices described in this notice and give you a copy of it.
- We will not use or share your information other than as described here unless you tell us we can in writing. If you tell us we can, you may change your mind at any time. Let us know in writing if you change your mind.

For more information see: www.hhs.gov/ocr/privacy/hipaa/understanding/consumers/noticepp.html.

Changes to the Terms of this Notice

We can change the terms of this notice, and the changes will apply to all information we have about you. The new notice will be available upon request, in our office, and on our web site.

This Addendum describes State laws governing the Pharmacy's use of your health information. The Pharmacy will comply with the applicable State law instead of the HIPAA Privacy Rule to the extent the State law provides greater privacy protection, or gives you more rights regarding your health information, than does the HIPAA Privacy Rule.

All States

The Pharmacy will not disclose the following categories of individual identifiable health information except in situations where the subject of the information has provided us with a written authorization allowing the release, or where we are permitted or required by state or federal law to make the disclosure. These categories include HIV/AIDS, mental health, communicable disease, alcohol and substance abuse, genetic information, information pertaining to minors and reproductive rights.

If you have questions or would like more information about Southeastern CrossCare's privacy policy, contact our Privacy Officer at 8000 Practitioner Parkway, Building 200, Jacksonville, Florida 32256 or toll

Walgreens

#05382 4330 HIGHWAY 22
MANDEVILLE, LA 70471
985-674-2551

905 3862 0041 07/03/2020 10:32 AM

FSA RX 2306184 B 0.00

TOTAL 0.00
CASH 0.00

THANK YOU FOR SHOPPING AT WALGREENS

YOU COULD HAVE EARNED AN ADDITIONAL 100
POINTS BY USING YOUR BALANCE REWARDS
CARD TODAY. RESTRICTIONS APPLY. FOR
TERMS AND CONDITIONS, VISIT
WALGREENS.COM/BALANCE.

NOT A MEMBER? JOIN NOW AT ANY REGISTER,
OR GO TO WALGREENS.COM/BALANCE.
ENROLLING IS QUICK, EASY AND FREE!
REDEEM POINTS FOR DOLLARS OFF FUTURE
PURCHASES.

RFN# 0538-2413-8621-2007-0303



SEE THE IMPACT OUR BACK TO SCHOOL ME TO
WE PROUD SUPPORTERS ARE MAKING ON
TEACHERS ACROSS AMERICA.

TRACK YOUR IMPACT:
WWW.WALGREENS.COM/METOWE

How are we doing?
Enter our monthly sweepstakes for
\$3,000 cash

Visit
WWW.WALGREENSLISTENS.COM

or call toll free
1-800-219-7451
within 72 hours to take a short
survey about this Walgreens visit

SURVEY#
0538-2413-862

PASSWORD
1200-7030-326


For contest rules, see store or
WWW.WALGREENSLISTENS.COM

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

| | | | | |
|-------------------|--|--------------------------|-----------------------------|---|
| PATIENT | JEFFREY BODIN | DOCTOR | K. MCLAUGHLIN, MD | DRUG DESCRIPTION |
| BIRTH DATE | 05/22/97 | | |  |
| MEDICATION | D-AMPHETAMINE SALT COMBO 30MG TABS | PATIENT ALLERGIES | <ALLERGY HISTORY NOT KNOWN> | |
| QUANTITY | 90 | | | |
| DIRECTIONS | TAKE ONE TABLET BY MOUTH THREE TIMES DAILY | | | PEACH |
| | | | | FRONT: b / 974 |
| | | | | BACK: 30 |

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablelets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. **TELL YOUR DOCTOR:** If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, ibuprofen or like products, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A severe and sometimes deadly problem called serotonin syndrome

may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable. Headache. Trouble sleeping. Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 828188 12T

JEFFREY BODIN
528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2306184-05382 DATE: 07/03/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC:00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD PLAN: ABHPA
MFG:TEVA GROUP# RX8834
DSB/JLC/JLC/JLC/JLC CLAIM REF# 201853776033212999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN
528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2306184-05382 DATE: 07/03/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC:00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD PLAN: ABHPA
MFG:TEVA GROUP# RX8834
DSB/JLC/JLC/JLC/JLC CLAIM REF# 201853776033212999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

WAITING

FRI 10:43AM

New


D-AMPHETAMINE SALT COMBO 30MG TABS

00555-0974-02

SAFE

QTY 90

20 DRAM



PEACH

FRONT: b / 974

BACK: 30

DSB/JLC/JLC/JLC/JLC

Med Guide

Walgreens

#05382 4330 HIGHWAY 22
MANDEVILLE, LA 70471
985-674-2551

810 8291 0041 08/03/2020 11:31 AM

FSA RX 2316069 B 0.00

TOTAL 0.00

CASH 0.00

THANK YOU FOR SHOPPING AT WALGREENS

GET MORE WITH BALANCE REWARDS,
REDEEM POINTS FOR SOMETHING EXTRA
IN A FUTURE PURCHASE. RESTRICTIONS
APPLY. FOR FULL DETAILS SEE
WALGREENS.COM/BALANCE.

RFN# 0538-2418-2919-2008-0303





POINT BALANCE 100

BALANCE REWARDS ACCT # *****0353

OPENING BALANCE 0

EVERYDAY POINTS - RX 100

CLOSING BALANCE 100

SEE THE IMPACT OUR BACK TO SCHOOL ME TO
WE PROUD SUPPORTERS ARE MAKING ON
TEACHERS ACROSS AMERICA.

TRACK YOUR IMPACT:
WWW.WALGREENS.COM/METOWE

Track Your Impact
See how together with our WE Teachers
Program Proud Supporters we are making a
positive impact on teachers in our local
communities: Walgreens.com/WeTeachers

How are we doing?
Enter our monthly sweepstakes for
\$3,000 cash

Visit
WWW.WALGREENSLISTENS.COM

or call toll free
1-800-219-7451

within 72 hours to take a short
survey about this Walgreens visit

SURVEY#
0538-2418-291

PASSWORD
9200-8030-326


For contest rules, see store or
WWW.WALGREENSLISTENS.COM

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

| | | |
|--|--|---|
| PATIENT JEFFREY BODIN | DOCTOR K. MCLAUGHLIN, MD | DRUG DESCRIPTION |
| BIRTH DATE 05/22/97 | |  |
| MEDICATION D-AMPHETAMINE SALT COMBO 30MG TABS | | |
| QUANTITY 90 | PATIENT ALLERGIES < ALLERGY HISTORY NOT KNOWN > | |
| DIRECTIONS TAKE 1 TABLET BY MOUTH THREE TIMES DAILY | | PEACH |
| | | FRONT: b / 974 |
| | | BACK: 30 |

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. TELL YOUR DOCTOR: If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. TELL YOUR DOCTOR: If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. TELL YOUR DOCTOR: If you have ever had any of these health problems: Drug abuse or stroke. TELL YOUR DOCTOR: If you have kidney disease. TELL YOUR DOCTOR: If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. TELL YOUR DOCTOR: If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. TELL YOUR DOCTOR: If you are taking any of these drugs: Linezolid or methylene blue. TELL YOUR DOCTOR: If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. WHAT DO I DO IF I MISS A DOSE? Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, ibuprofen or like products, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A severe and sometimes deadly problem called serotonin syndrome

may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable. Headache. Trouble sleeping. Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188 12T

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2316069-05382

DATE: 08/03/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD
MFG: TEVA
MOJ/LSR/LSR/LSR/JLC

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 202164112643115999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2316069-05382

DATE: 08/03/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD
MFG: TEVA
MOJ/LSR/LSR/LSR/JLC

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 202164112643115999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

WAITING

MON 11:39AM

New


D-AMPHETAMINE SALT COMBO 30MG TABS

00555-0974-02

SAFE

QTY 90

20 DRAM



PEACH

FRONT: b / 974

BACK: 30

MOJ/LSR/LSR/LSR/JLC

Med Guide

Mandeville Pharmacy
4655 Highway 190
Mandeville, LA 70448-3582
Phone: (985) 778-2306
Fax: (985) 778-2404

Aug 5 2020 9:57AM
Receipt # 38347
Cashier: POORNG C
Drawer #: POS01 - 3275

Rx 43691-00 - JEFFREY 72.99 F
BODIN

Subtotal 72.99
Tax Total 3.57

Items: 1 Total 76.56

Visa Tendered 76.56
Acct #...1006
Approval #...025710

F = FLEX/FSA/HRA Eligible
FSA Eligible \$76.56

Signature required for the following:

- Payment(s) require a signature
- Receipt of Prescriptions
- Receipt of Privacy Practices

**Impacted By
COVID-19**

Signature Captured Electronically

I agree to pay above total amount according to
card issuer agreement.

MID: 9139901
Entry: TAP



Scan This QR Code for Patient Specific
Medication Information

"Where the Customer is King/Queen"




N0043691-00-00
JEFFREY BODIN
RXLOC
Price 72.99
08/05/2020
Sig: _____

MA

N0043691 08/05/2020
JEFFREY BODIN
Rej: _____ N/A: _____

MALCOLM ARMSTRONG
DEXTROAMP-AMPHETAMIN 10 MG TA

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306
JEFFREY BODIN
528 BEAU CHENE DR (985) 272-8989
MANDEVILLE, LA 70471-1777 PP: RXLO Dr. Kevin McLaughlin
Rx: N0043691 00 08/05/2020 **\$72.99** Rph: MA
DEXTROAMP-AMPHETAMIN 10 MG TAB 68382-0952-01 Qty 90



Mandeville Pharmacy 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

RECEIPT



Rx: N0043691 00
JEFFREY BODIN
528 BEAU CHENE DR

08/05/2020 (985) 272-8989

DOB: 05/22/1997

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

Auth:

PP: RXLOC

NO REFILLS

Rph: MALCOLM ARMSTRONG
(985) 867-5494
68382-0952-01
PHCY ID: 1548616998

Copay

\$72.99

MANDEVILLE PHARMACY

1655 HIGHWAY 190
MANDEVILLE, LA 70448-3582

(985) 778-2306

08/05/2020

Rx: N0043691 00
JEFFREY BODIN

528 BEAU CHENE DR
MANDEVILLE, LA 70471-1777

(985) 272-8989

Dr. Kevin McLaughlin
DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:
NO REFILLS

Rph: MALCOLM ARMSTRONG
(985) 867-5494
68382-0952-01 Qty 90
PHCY ID: 1548616998

**DUPLICATE
RECEIPT**

Copay **\$72.99**
PP: RXLOC

Call your Doctor for medical advice about side effects.
You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"

PRESCRIPTION NOTES

Mandeville Pharmacy
1655 Highway 190
Mandeville, LA 70448-3582

For: JEFFREY BODIN
Rx: 43691
Date: August 05, 2020
Message ID: 81842656-407

MEDICATION GUIDE **Adderall® (ADD-ur-all) CII** **(Dextroamphetamine Saccharate, Amphetamine Aspartate, Dextroamphetamine Sulfate and Amphetamine Sulfate Tablets)**

Rx only

Read the Medication Guide that comes with Adderall® before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with Adderall®.

What is the most important information I should know about Adderall®?

The following have been reported with use of Adderall® and other stimulant medicines.

1. Heart-Related problems:

- sudden death in patients who have heart problems or heart defects
- stroke and heart attack in adults
- increased blood pressure and heart rate

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting Adderall®.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with Adderall®.

Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Adderall®.

2. Mental (Psychiatric) problems:

All Patients

- new or worse behavior and thought problems
- new or worse bipolar illness
- new or worse aggressive behavior or hostility

Children and Teenagers

- new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking Adderall®, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

3. Circulation problems in fingers and toes [Peripheral vasculopathy, including Raynaud's phenomenon]:

- fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes.

Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking Adderall®.

What is Adderall®?

Adderall® is a central nervous system stimulant prescription medicine. **It is used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD).** Adderall® may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD. Adderall® should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Adderall® is also used in the treatment of a sleep disorder called narcolepsy.

Adderall® is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep Adderall® in a safe place to prevent misuse and abuse. Selling or giving away Adderall® may harm others, and is against the law.

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take Adderall®?

Adderall® should not be taken if you or your child:

- have heart disease or hardening of the arteries
 - have moderate to severe high blood pressure
 - have hyperthyroidism
 - have an eye problem called glaucoma
 - are very anxious, tense, or agitated
 - have a history of drug abuse
 - are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOI.
 - are sensitive to, allergic to, or had a reaction to other stimulant medicines
- Adderall® is not recommended for use in children less than 3 years old.

Adderall® may not be right for you or your child. Before starting Adderall® tell your or your child's doctor about all health conditions (or a family history of) including:

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- liver or kidney problems

- thyroid problems
- seizures or have had an abnormal brain wave test (EEG)
- circulation problems in fingers or toes

Tell your doctor if you or your child are pregnant, planning to become pregnant, or breastfeeding.

Can Adderall® be taken with other medicines?

Tell your doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements. Adderall® and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking Adderall®.

Your doctor will decide whether Adderall® can be taken with other medicines.

Especially tell your doctor if you or your child take:

- anti-depression medicines including MAOIs
- blood pressure medicines
- seizure medicines
- blood thinner medicines
- cold or allergy medicines that contain decongestants
- stomach acid medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking Adderall® without talking to your doctor first.

How should Adderall® be taken?

- **Take Adderall® exactly as prescribed.** Your doctor may adjust the dose until it is right for you or your child.
- Adderall® tablets are usually taken two to three times a day. The first dose is usually taken when you first wake in the morning. One or two more doses may be taken during the day, 4 to 6 hours apart.
- Adderall® can be taken with or without food.
- From time to time, your doctor may stop Adderall® treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure while taking Adderall®. Children should have their height and weight checked often while taking Adderall®. Adderall® treatment may be stopped if a problem is found during these check-ups.
- **If you or your child take too much Adderall® or overdoses, call your doctor or poison control center right away, or get emergency treatment.**

What are possible side effects of Adderall®?

See **“What is the most important information I should know about Adderall®?”** for information on reported heart and mental problems.

Other serious side effects include:

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision

Common side effects include:

- stomach ache
- decreased appetite
- nervousness

Adderall® may affect your or your child’s ability to drive or do other dangerous activities.

Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist

for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Adderall®?

- Store Adderall® in a safe place at room temperature, 20° to 25°C (68° to 77°F).
- **Keep Adderall® and all medicines out of the reach of children.**

General information about Adderall®

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Adderall® for a condition for which it was not prescribed. Do not give Adderall® to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about Adderall®. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Adderall® that was written for healthcare professionals. For more information about Adderall®, please contact Teva Pharmaceuticals at 1-888-838-2872.

What are the ingredients in Adderall®?

Active Ingredient: dextroamphetamine saccharate, amphetamine aspartate monohydrate, dextroamphetamine sulfate, USP, amphetamine sulfate, USP

Inactive Ingredients: lactitol, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, and other ingredients.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

[Teva Select Brands, Horsham, PA 19044](#)

Division of Teva Pharmaceuticals USA

Revised October 2015

Patient Education - Dextroamp-Amphetamin 10 Mg Tab

CONSUMER INFORMATION USE AND DISCLAIMER This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Dextroamphetamine and Amphetamine Tablets

PRONUNCIATION (deks troe am FET a meen & am FET a meen)

BRAND NAME(S): Adderall

WARNING This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

WHAT IS THIS DRUG USED FOR? It is used to treat attention deficit problems with hyperactivity.; It is used to treat narcolepsy.; It may be given to you for other reasons. Talk with the doctor.

WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.; If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics.; If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid.; If you have ever had any of these health problems: Drug abuse or stroke.; If you have kidney disease.; If you are taking any of these drugs: Acetazolamide or sodium bicarbonate.; If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen.; If you are taking any of these drugs: Linezolid or methylene blue.; If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

WHAT ARE SOME THINGS I NEED TO KNOW OR DO WHILE I TAKE THIS DRUG? Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. ; Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.; If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered.; If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug.; You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor.; This drug may cause high blood pressure.; Check blood pressure and heart rate as the doctor has told you.; Have blood work checked as you have been told by the doctor. Talk with the doctor.; This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.; Do not take antacids with this drug.; Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.; New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life.; A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache.; This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.; Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child.; Talk with your doctor before you drink alcohol.; Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.; Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.; Not able to get or keep an erection.; Change in sex interest.; Seizures.; Trouble controlling body movements.; Restlessness.; Change in eyesight.; For males, erections (hard penis) that happen often or that last a long time.; Change in color of hands or feet from pale to blue or red.; Numbness, pain, tingling, or cold feeling of the hands or feet.; Any sores or wounds on the fingers or toes.; Muscle pain or weakness, dark urine, or trouble passing urine.; Pain when passing urine.; Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out.

WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable.; Headache.; Trouble sleeping.; Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry.; Feeling dizzy, tired, or weak.; Dry mouth.; Bad taste in your mouth.; Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.; Take with or without food.; Take last dose of the day at least 4 hours before bedtime.

WHAT DO I DO IF I MISS A DOSE? Take a missed dose as soon as you think about it.; If it is close to the time for your next dose, skip the missed dose and go back to your normal time.; Do not take 2 doses at the same time or extra doses.

HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.; Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.; Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

GENERAL DRUG FACTS If your symptoms or health problems do not get better or if they become worse, call your doctor. ; Do not share your drugs with others and do not take anyone else's drugs.; This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If

you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.; If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Copyright 2020 UpToDate, Inc. All Rights Reserved. (Issue 20.3.3.005)

Mandeville Pharmacy
1655 Highway 190
Mandeville, LA 70448-3582
Phone: (985) 778-2306
Fax: (985) 778-2404

Sep 5 2020 10:34AM
Receipt # 39457
Cashier: NORMAN S
Drawer #: POS01 - 3342

Rx 44994-00 - JEFFREY BODIN 72.99 F

Subtotal 72.99
Tax Total 3.57

Items: 1 Total 76.56

Visa Tendered 76.56
Acct #...1006
Approval #...033411
Trans ID#...488670

F = FLEX/FSA/HRA Eligible
FSA Eligible \$76.56

Signature required for the following:

- Receipt of Prescriptions
- Receipt of Privacy Practices

**Impacted By
COVID-19**

Signature Captured Electronically

I agree to pay above total amount according to card issuer agreement.

MID: 9139901
Term ID: 1790344
App: US DEBIT
AID: A0000000980840
AC: ARQC B80E9D249AD25042
Entry: INSERT
CVM: Pin Verified
DebitSale



Scan This QR Code for Patient Specific Medication Information

"Where the Customer is King/Queen"




N0044994-00-00
JEFFREY BODIN
RXLOC
Price 72.99
09/05/2020
Sig: _____

NS

N0044994 09/05/2020
JEFFREY BODIN
Rej: _____ N/A: _____

NORMAN SMITH
DEXTROAMP-AMPHETAMIN 10 MG TAB

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306
JEFFREY BODIN
528 BEAU CHENE DR (985) 272-8989
MANDEVILLE, LA 70471-1777 PP: RXLO Dr. Kevin McLaughlin
Rx: N0044994 00 09/05/2020 **\$72.99** Rph: NS
DEXTROAMP-AMPHETAMIN 10 MG TAB 00555-0972-02 Qty 90



Mandeville Pharmacy 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

Rx: N0044994 00
JEFFREY BODIN
528 BEAU CHENE DR

09/05/2020 (985) 272-8989

RECEIPT



DOB: 05/22/1997

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

Auth:

PP: **RXLOC**

NO REFILLS

Rph: NORMAN SMITH

(985) 867-5494

00555-0972-02

PHCY ID: 1548616998

Copay

\$72.99

MANDEVILLE PHARMACY

1655 HIGHWAY 190
MANDEVILLE, LA 70448-3582

(985) 778-2306

09/05/2020

Rx: N0044994 00
JEFFREY BODIN

528 BEAU CHENE DR
MANDEVILLE, LA 70471-1777

(985) 272-8989

Dr. Kevin McLaughlin
DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:
NO REFILLS

Rph: NORMAN SMITH

(985) 867-5494

00555-0972-02

Qty 90

PHCY ID: 1548616998

**DUPLICATE
RECEIPT**

Copay **\$72.99**

PP: **RXLOC**

Call your Doctor for medical advice about side effects.
You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"

PRESCRIPTION NOTES

Mandeville Pharmacy
1655 Highway 190
Mandeville, LA 70448-3582

For: JEFFREY BODIN
Rx: 44994
Date: September 05, 2020
Message ID: 38868793-407

MEDICATION GUIDE

Adderall[®] (ADD-ur-all) CII

(Dextroamphetamine Saccharate, Amphetamine Aspartate, Dextroamphetamine Sulfate and Amphetamine Sulfate Tablets)

Rx only

Read the Medication Guide that comes with Adderall[®] before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with Adderall[®].

What is the most important information I should know about Adderall[®]?

The following have been reported with use of Adderall[®] and other stimulant medicines.

1. Heart-Related problems:

- sudden death in patients who have heart problems or heart defects
- stroke and heart attack in adults
- increased blood pressure and heart rate

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting Adderall[®].

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with Adderall[®].

Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Adderall[®].

2. Mental (Psychiatric) problems:

All Patients

- new or worse behavior and thought problems
- new or worse bipolar illness
- new or worse aggressive behavior or hostility

Children and Teenagers

- new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking Adderall[®], especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

3. Circulation problems in fingers and toes [Peripheral vasculopathy, including Raynaud's phenomenon]:

- fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes.

Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking Adderall[®].

What is Adderall[®]?

Adderall[®] is a central nervous system stimulant prescription medicine. **It is used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD).** Adderall[®] may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD. Adderall[®] should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Adderall[®] is also used in the treatment of a sleep disorder called narcolepsy.

Adderall[®] is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep Adderall[®] in a safe place to prevent misuse and abuse. Selling or giving away Adderall[®] may harm others, and is against the law.

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take Adderall[®]?

Adderall[®] should not be taken if you or your child:

- have heart disease or hardening of the arteries
 - have moderate to severe high blood pressure
 - have hyperthyroidism
 - have an eye problem called glaucoma
 - are very anxious, tense, or agitated
 - have a history of drug abuse
 - are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOI.
 - are sensitive to, allergic to, or had a reaction to other stimulant medicines
- Adderall[®] is not recommended for use in children less than 3 years old.

Adderall[®] may not be right for you or your child. Before starting Adderall[®] tell your or your child's doctor about all health conditions (or a family history of) including:

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- liver or kidney problems

- thyroid problems
- seizures or have had an abnormal brain wave test (EEG)
- circulation problems in fingers or toes

Tell your doctor if you or your child are pregnant, planning to become pregnant, or breastfeeding.

Can Adderall® be taken with other medicines?

Tell your doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements. Adderall® and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking Adderall®.

Your doctor will decide whether Adderall® can be taken with other medicines.

Especially tell your doctor if you or your child take:

- anti-depression medicines including MAOIs
- blood pressure medicines
- seizure medicines
- blood thinner medicines
- cold or allergy medicines that contain decongestants
- stomach acid medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking Adderall® without talking to your doctor first.

How should Adderall® be taken?

- **Take Adderall® exactly as prescribed.** Your doctor may adjust the dose until it is right for you or your child.
- Adderall® tablets are usually taken two to three times a day. The first dose is usually taken when you first wake in the morning. One or two more doses may be taken during the day, 4 to 6 hours apart.
- Adderall® can be taken with or without food.
- From time to time, your doctor may stop Adderall® treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure while taking Adderall®. Children should have their height and weight checked often while taking Adderall®. Adderall® treatment may be stopped if a problem is found during these check-ups.
- **If you or your child take too much Adderall® or overdoses, call your doctor or poison control center right away, or get emergency treatment.**

What are possible side effects of Adderall®?

See **“What is the most important information I should know about Adderall®?”** for information on reported heart and mental problems.

Other serious side effects include:

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision

Common side effects include:

- stomach ache
- decreased appetite
- nervousness

Adderall® may affect your or your child's ability to drive or do other dangerous activities.

Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist

for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Adderall®?

- Store Adderall® in a safe place at room temperature, 20° to 25°C (68° to 77°F).
- **Keep Adderall® and all medicines out of the reach of children.**

General information about Adderall®

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Adderall® for a condition for which it was not prescribed. Do not give Adderall® to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about Adderall®. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Adderall® that was written for healthcare professionals. For more information about Adderall®, please contact Teva Pharmaceuticals at 1-888-838-2872.

What are the ingredients in Adderall®?

Active Ingredient: dextroamphetamine saccharate, amphetamine aspartate monohydrate, dextroamphetamine sulfate, USP, amphetamine sulfate, USP

Inactive Ingredients: lactitol, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, and other ingredients.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Teva Select Brands, Horsham, PA 19044

Division of Teva Pharmaceuticals USA

Revised October 2015

Walgreens

#05382 4330 HIGHWAY 22
MANDEVILLE, LA 70471
985-674-2551

878 3097 0041 09/05/2020 11:02 AM

FSA RX 2327527 B 0.00

TOTAL 0.00

CASH 0.00

THANK YOU FOR SHOPPING AT WALGREENS

GET MORE WITH BALANCE REWARDS,
REDEEM POINTS FOR SOMETHING EXTRA
IN A FUTURE PURCHASE. RESTRICTIONS
APPLY. FOR FULL DETAILS SEE
WALGREENS.COM/BALANCE.

RFN# 0538-2413-0975-2009-0503





POINT BALANCE 600

BALANCE REWARDS ACCT # *****8423

OPENING BALANCE 500

EVERYDAY POINTS - RX 100

CLOSING BALANCE 600

SEE THE IMPACT OUR BACK TO SCHOOL ME TO
WE PROUD SUPPORTERS ARE MAKING ON
TEACHERS ACROSS AMERICA.

TRACK YOUR IMPACT:
WWW.WALGREENS.COM/METOWE

Track Your Impact

See how together with our WE Teachers
Program Proud Supporters we are making a
positive impact on teachers in our local
communities: Walgreens.com/WeTeachers

How are we doing?
Enter our monthly sweepstakes for
\$3,000 cash

Visit
WWW.WALGREENSLISTENS.COM

or call toll free
1-800-219-7451

within 72 hours to take a short
survey about this Walgreens visit

SURVEY#
0538-2413-097

PASSWORD
5200-9050-326


For contest rules, see store or
WWW.WALGREENSLISTENS.COM

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

| | | |
|--|--|---|
| PATIENT JEFFREY BODIN | DOCTOR K. MCLAUGHLIN, MD | DRUG DESCRIPTION |
| BIRTH DATE 05/22/97 | |  |
| MEDICATION D-AMPHETAMINE SALT COMBO 30MG TABS | PATIENT ALLERGIES < ALLERGY HISTORY NOT KNOWN > | |
| QUANTITY 90 | | PEACH |
| DIRECTIONS TAKE 1 TABLET BY MOUTH THREE TIMES DAILY | | FRONT: b / 974 BACK: 30 |

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. **TELL YOUR DOCTOR:** If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in

life. A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable. Headache. Trouble sleeping. Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188 12T

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2327527-05382

DATE: 09/05/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD
MFG: TEVA
MQB/VNV/VNV/VNV/VNV

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 202493906348177999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2327527-05382

DATE: 09/05/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD
MFG: TEVA
MQB/VNV/VNV/VNV/VNV

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 202493906348177999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt


Pharmacy use only

WAITING

SAT 11:05AM
New

D-AMPHETAMINE SALT COMBO 30MG TABS
00555-0974-02
SAFE

QTY 90
20 DRAM



PEACH
FRONT: b / 974
BACK: 30

MQB/VNV/VNV/VNV/VNV

Med Guide

Mandeville Pharmacy
1655 Highway 190
Mandeville, LA 70448-3582
Phone: (985) 778-2306
Fax: (985) 778-2404

Oct 6 2020 5:08PM
Receipt # 40556
Cashier: NORMAN S
Drawer #: POS01 - 3405

Rx 46231-00 - JEFFREY 72.99 F
BODIN

Subtotal 72.99
Tax Total 3.57

Items: 1 Total 76.56

Visa Tendered 76.56
Acct #...1006
Approval #...070818
Trans ID#...500390

F = FLEX/FSA/HRA Eligible
FSA Eligible \$76.56

Signature required for the following:

- Receipt of Prescriptions
- Receipt of Privacy Practices

**Impacted By
COVID-19**

Signature Captured Electronically

I agree to pay above total amount according to
card issuer agreement.

MID: 9139901
Term ID: 1790344
App: US DEBIT
AID: A0000000980840
AC: ARQC FE419CBF0B765051
Entry: INSERT
CVM: Pin Verified
DebitSale



Scan This QR Code for Patient Specific
Medication Information

"Where the Customer is King/Queen"




N0046231-00-00
JEFFREY BODIN
RXLOC
Price 72.99
10/06/2020
Sig: _____

NS

N0046231 10/06/2020
JEFFREY BODIN
Rej: _____ N/A: _____

NORMAN SMITH
DEXTROAMP-AMPHETAMIN 10 MG T/

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306
JEFFREY BODIN
528 BEAU CHENE DR (985) 272-8989
MANDEVILLE, LA 70471-1777 PP: RXLOC Dr. Kevin McLaughlin
Rx: N0046231 00 10/06/2020 **\$72.99** Rph: NS
DEXTROAMP-AMPHETAMIN 10 MG TAB 68382-0952-01 Qty 90

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

Rx: N0046231 00
JEFFREY BODIN
528 BEAU CHENE DR

10/06/2020 (985) 272-8989

RECEIPT



DOB: 05/22/1997

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

Auth:

PP: RXLOC

NO REFILLS

Rph: NORMAN SMITH
(985) 867-5494
68382-0952-01
PHCY ID: 1548616998

Copay \$72.99

*ICU #35
→ Thursday*

MANDEVILLE PHARMACY

1655 HIGHWAY 190
MANDEVILLE, LA 70448-3582

(985) 778-2306

10/06/2020

Rx: N0046231 00
JEFFREY BODIN

528 BEAU CHENE DR

(985) 272-8989

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:

NO REFILLS

Rph: NORMAN SMITH

(985) 867-5494

68382-0952-01

Qty 90

PHCY ID: 1548616998

**DUPLICATE
RECEIPT**

Copay \$72.99

PP: RXLOC

Call your Doctor for medical advice about side effects.
You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"


PRESCRIPTION NOTES

N0046231-00-00
JEFFREY BODIN
RXLOC
Price 72.99
10/06/2020
Sig: _____


NS

N0046231 10/06/2020
JEFFREY BODIN
Rej: _____ N/A: _____

NORMAN SMITH
DEXTROAMP-AMPHETAMIN 10 MG T/

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

JEFFREY BODIN (985) 272-8989
528 BEAU CHENE DR
MANDEVILLE, LA 70471-1777 PP: RXLO Dr. Kevin McLaughlin
Rx: N0046231 00 10/06/2020 **\$72.99** Rph: NS
DEXTROAMP-AMPHETAMIN 10 MG TAB 68382-0952-01 Qty 90



 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

Rx: N0046231 00
JEFFREY BODIN
528 BEAU CHENE DR

10/06/2020 (985) 272-8989

RECEIPT



DOB: 05/22/1997

*#35 completion 200
Done*

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

PP: RXLOC

Auth:

NO REFILLS

Rph: NORMAN SMITH
(985) 867-5494
68382-0952-01
PHCY ID: 1548616998

Copay

N
\$72.99
C

MANDEVILLE PHARMACY

1655 HIGHWAY 190
MANDEVILLE, LA 70448-3582

(985) 778-2306

10/06/2020

Rx: N0046231 00
JEFFREY BODIN

528 BEAU CHENE DR
MANDEVILLE, LA 70471-1777

(985) 272-8989

Dr. Kevin McLaughlin
DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:
NO REFILLS

Rph: NORMAN SMITH
(985) 867-5494
68382-0952-01 Qty 90
PHCY ID: 1548616998

**DUPLICATE
RECEIPT**

Copay **\$72.99**
PP: RXLOC

Call your Doctor for medical advice about side effects.
You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"

PRESCRIPTION NOTES

Walgreens

#09990 71041 HIGHWAY 21
COVINGTON, LA 70433
985-875-0715

148 2138 0022 10/05/2020 12:02 PM

ESSIE GL NL MAKEK THE CUT .460Z
09500802125 A 11.50
RETURN VALUE 11.50
ESSIE GL CTR CLR FARY TALOR .460Z
09500802098 A 11.50
RETURN VALUE 11.50
~~ACT RESTOR M/W C/SPLSH MINT 330Z~~
04116709806 OTC A ~~7.49~~

SUBTOTAL 30.49
SALES TAX A=8.7% 2.65
TOTAL 33.14
AMEX ACCT 7025 33.14
CHANGE .00

AID A000000025010801
AMERICAN EXPRESS
Integrated chip card

THANK YOU FOR SHOPPING AT WALGREENS

GET MORE WITH BALANCE REWARDS,
REDEEM POINTS FOR SOMETHING EXTRA
IN A FUTURE PURCHASE. RESTRICTIONS
APPLY. FOR FULL DETAILS SEE
WALGREENS.COM/BALANCE.

RFN# 0999-0222-1380-2010-0503





POINT BALANCE 100

BALANCE REWARDS ACCT # *****0750

How are we doing?
Enter our monthly sweepstakes for
\$3,000 cash

Visit
WWW.WALGREENSLISTENS.COM

or call toll free
1-800-219-7451
within 72 hours to take a short
survey about this Walgreens visit

SURVEY#
0999-0222-138

PASSWORD
0201-0050-326


For contest rules, see store or
WWW.WALGREENSLISTENS.COM

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

| | | | | |
|-------------------|--|--------------------------|-------------------------------|---|
| PATIENT | JEFFREY BODIN | DOCTOR | K. MCLAUGHLIN, MD | DRUG DESCRIPTION |
| BIRTH DATE | 05/22/97 | | |  |
| MEDICATION | D-AMPHETAMINE SALT COMBO 30MG TABS | PATIENT ALLERGIES | < ALLERGY HISTORY NOT KNOWN > | |
| QUANTITY | 90 | | | PEACH |
| DIRECTIONS | TAKE 1 TABLET BY MOUTH THREE TIMES DAILY | | | FRONT: b / 974 |
| | | | | BACK: 30 |

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. **TELL YOUR DOCTOR:** If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in

life. A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excited. Headache. Trouble sleeping. Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188

JEFFREY BODIN
528 Beau Chene Dr, Mandeville, LA 70471777
(985)272-8989
RX # 2338451-05382 DATE: 10/06/20

D-AMPHETAMINE SALT COMBO 30MG TABS
QTY: 90 NO REFILLS
New NDC: 00555-0974-02 \$ 0.00

K. MCLAUGHLIN, MD PLAN: ABHPA
MFG: TEVA GROUP# RX8834
AAT/LSR/LSR/LSR/JLC CLAIM REF# 202804272761021999

Walgreens 4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551 **Customer Receipt**

JEFFREY BODIN
528 Beau Chene Dr, Mandeville, LA 70471777
(985)272-8989
RX # 2338451-05382 DATE: 10/06/20

D-AMPHETAMINE SALT COMBO 30MG TABS
QTY: 90 NO REFILLS
New NDC: 00555-0974-02 \$ 0.00

K. MCLAUGHLIN, MD PLAN: ABHPA
MFG: TEVA GROUP# RX8834
AAT/LSR/LSR/LSR/JLC CLAIM REF# 202804272761021999


Walgreens 4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551 **Duplicate Receipt**

Pharmacy use only

WAITING
TUE 12:06PM
New

D-AMPHETAMINE SALT COMBO 30MG TABS
00555-0974-02
SAFE

QTY 90
20 DRAM



PEACH
FRONT: b / 974
BACK: 30

AAT/LSR/LSR/LSR/JLC

Med Guide

MEDICATION GUIDE

Adderall® (ADD-ur-all) (CII)

(Dextroamphetamine Saccharate, Amphetamine Aspartate, Dextroamphetamine Sulfate and Amphetamine Sulfate Tablets)

Rx only

Read the Medication Guide that comes with Adderall® before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with Adderall®.

What is the most important information I should know about Adderall®?

The following have been reported with use of Adderall® and other stimulant medicines.

1. Heart-related problems:

- sudden death in patients who have heart problems or heart defects
- stroke and heart attack in adults
- increased blood pressure and heart rate

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting Adderall®.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with Adderall®.

Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Adderall®.

2. Mental (Psychiatric) problems:

All Patients

- new or worse behavior and thought problems
- new or worse bipolar illness
- new or worse aggressive behavior or hostility

Children and Teenagers

- new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking Adderall®, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

3. Circulation problems in fingers and toes [Peripheral vasculopathy, including Raynaud's phenomenon]:

- fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in your fingers or toes.

Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking Adderall®.

What is Adderall®?

Adderall® is a central nervous system stimulant prescription medicine. **It is used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD).** Adderall® may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD.

Adderall® should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Adderall® is also used in the treatment of a sleep disorder called narcolepsy.

Adderall® is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep Adderall® in a safe place to prevent misuse and abuse. Selling or giving away Adderall® may harm others, and is against the law.

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take Adderall®?

Adderall® should not be taken if you or your child:

- have heart disease or hardening of the arteries
- have moderate to severe high blood pressure
- have hyperthyroidism
- have an eye problem called glaucoma
- are very anxious, tense, or agitated
- have a history of drug abuse
- are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOI.
- are sensitive to, allergic to, or had a reaction to other stimulant medicines

Adderall® is not recommended for use in children less than 3 years old.

Adderall® may not be right for you or your child. Before starting Adderall® tell your or your child's doctor about all health conditions (or a family history of) including:

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- liver or kidney problems
- thyroid problems
- seizures or have had an abnormal brain wave test (EEG)
- circulation problems in fingers and toes

Tell your doctor if you or your child are pregnant, planning to become pregnant, or breastfeeding.

Can Adderall® be taken with other medicines?

Tell your doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements. Adderall® and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking Adderall®.

Your doctor will decide whether Adderall® can be taken with other medicines.

Especially tell your doctor if you or your child take:

- anti-depression medicines including MAOIs
- blood pressure medicines
- seizure medicines
- blood thinner medicines
- cold or allergy medicines that contain decongestants
- stomach acid medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking Adderall® without talking to your doctor first.

How should Adderall® be taken?

- **Take Adderall® exactly as prescribed.** Your doctor may adjust the dose until it is right for you or your child.
- Adderall® tablets are usually taken two to three times a day. The first dose is usually taken when you first wake in the morning. One or two more doses may be taken during the day, 4 to 6 hours apart.
- Adderall® can be taken with or without food.
- From time to time, your doctor may stop Adderall® treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure while taking Adderall®. Children should have their height and weight checked often while taking Adderall®. Adderall® treatment may be stopped if a problem is found during these check-ups.
- **If you or your child take too much Adderall® or overdoses, call your doctor or poison control center right away, or get emergency treatment.**

What are possible side effects of Adderall®?

See “**What is the most important information I should know about Adderall®?**” for information on reported heart and mental problems.

Other serious side effects include:

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision
- Serotonin syndrome. A potentially life-threatening problem called serotonin syndrome can happen when medicines such as Adderall® are taken with certain other medicines. Symptoms of serotonin syndrome may include:
 - agitation, hallucinations, coma or other changes in mental status
 - problems controlling your movements or muscle twitching
 - fast heartbeat
 - high or low blood pressure
 - sweating or fever
 - nausea or vomiting
 - diarrhea
 - muscle stiffness or tightness

Common side effects include:

- stomach ache
- decreased appetite
- nervousness

Adderall® may affect your or your child’s ability to drive or do other dangerous activities. Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Adderall®?

- Store Adderall® in a safe place at room temperature, 20° to 25°C (68° to 77°F).
- **Keep Adderall® and all medicines out of the reach of children.**

General information about Adderall®

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Adderall® for a condition for which it was not prescribed. Do not give Adderall® to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about Adderall®. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Adderall® that was written for healthcare professionals. For more information about Adderall®, please contact Teva Pharmaceuticals at 1-888-838-2872.

What are the ingredients in Adderall®?

Active Ingredient: dextroamphetamine saccharate, amphetamine aspartate monohydrate, dextroamphetamine sulfate, USP, and amphetamine sulfate, USP

Inactive Ingredients: lactitol, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, and other ingredients.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Teva Select Brands, Horsham, PA 19044
Division of Teva Pharmaceuticals USA

Revised January 2017

Mandeville Pharmacy
1655 Highway 190
Mandeville, LA 70448-3582
Phone: (985) 778-2306
Fax: (985) 778-2404

Nov 6 2020 1:09PM
Receipt # 41649
Cashier: NORMAN S
Drawer #: POS01 - 3466

Rx 47416-00 - JEFFREY 72.99 F
BODIN

Subtotal 72.99

Tax Total 3.57

Items: 1 Total 76.56

Visa Tendered 76.56

Acct #...1006

Approval #...030914

Trans ID#...511920

F = FLEX/FSA/HRA Eligible
FSA Eligible \$76.56

Signature required for the following:

- Receipt of Prescriptions
- Receipt of Privacy Practices

**Impacted By
COVID-19**

Signature Captured Electronically

I agree to pay above total amount according to
card issuer agreement.

MID: 9139901

Term ID: 1790344

App: US DEBIT

AID: A0000000980840

AC: ARQC E7731A5B5D517FD7

Entry: INSERT

CVM: Pin Verified

DebitSale



Scan This QR Code for Patient Specific
Medication Information

"Where the Customer is King/Queen"





Mandeville Pharmacy 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

RECEIPT



Rx: N0047416 00
JEFFREY BODIN
 528 BEAU CHENE DR

11/06/2020 (985) 272-8989

DOB: 05/22/1997

TW \$50

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

Auth:

PP: **RXLOC**

NO REFILLS

Rph: NORMAN SMITH
 (985) 867-5494
 68382-0952-01
 PHCY ID: 1548616998

Copay

\$72.99

MANDEVILLE PHARMACY

1655 HIGHWAY 190
 MANDEVILLE, LA 70448-3582

(985) 778-2306

11/06/2020

Rx: N0047416 00
JEFFREY BODIN

528 BEAU CHENE DR
 MANDEVILLE, LA 70471-1777

(985) 272-8989

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:

NO REFILLS

Rph: NORMAN SMITH

(985) 867-5494

68382-0952-01

Qty 90

PHCY ID: 1548616998

**DUPLICATE
 RECEIPT**

Copay **\$72.99**

PP: **RXLOC**

Call your Doctor for medical advice about side effects.
 You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"


PRESCRIPTION NOTES

N0047416-00-00
JEFFREY BODIN
RXLOC
Price 72.99
11/06/2020
Sig: _____


NS

N0047416 11/06/2020
JEFFREY BODIN
Rej: _____ N/A: _____

NORMAN SMITH
DEXTROAMP-AMPHETAMIN 10 MG T/

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

JEFFREY BODIN
528 BEAU CHENE DR (985) 272-8989
MANDEVILLE, LA 70471-1777 PP: RXLO Dr. Kevin McLaughlin
Rx: N0047416 00 11/06/2020 **\$72.99** Rph: NS
DEXTROAMP-AMPHETAMIN 10 MG TAB 68382-0952-01 Qty 90



 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448
985-778-2306

Rx: N0047416 00
JEFFREY BODIN
528 BEAU CHENE DR

11/06/2020 (985) 272-8989

RECEIPT



MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

Auth:
NO REFILLS


PP: RXLOC

Rph: NORMAN SMITH
(985) 867-5494
68382-0952-01
PHCY ID: 1548616998

DOB: 05/22/1997

Completion 100 \$50

Copay

 **\$72.99**

MANDEVILLE PHARMACY

1655 HIGHWAY 190
MANDEVILLE, LA 70448-3582

(985) 778-2306

Rx: N0047416 00
JEFFREY BODIN

11/06/2020

528 BEAU CHENE DR
MANDEVILLE, LA 70471-1777

(985) 272-8989

Dr. Kevin McLaughlin
DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:
NO REFILLS

Rph: NORMAN SMITH
(985) 867-5494
68382-0952-01 Qty 90
PHCY ID: 1548616998

**DUPLICATE
RECEIPT**

Copay **\$72.99**
PP: RXLOC

Call your Doctor for medical advice about side effects.
You may report side effects to the FDA at: 1-800-FDA-1088.

We Appreciate Your Business!

"Where the Customer is King/Queen"

PRESCRIPTION NOTES

Walgreens

#05382 4330 HIGHWAY 22
MANDEVILLE, LA 70471
985-674-2551

841 1935 0041 11/06/2020 12:42 PM

FSA RX 2349453 B 0.00

TOTAL 0.00
CASH 0.00

THANK YOU FOR SHOPPING AT WALGREENS

GET MORE WITH BALANCE REWARDS,
REDEEM POINTS FOR SOMETHING EXTRA
IN A FUTURE PURCHASE. RESTRICTIONS
APPLY. FOR FULL DETAILS SEE
WALGREENS.COM/BALANCE.

RFN# 0538-2411-9352-2011-0603





POINT BALANCE 600

BALANCE REWARDS ACCT # *****8423

SEE THE IMPACT OUR BACK TO SCHOOL ME TO
WE PROUD SUPPORTERS ARE MAKING ON
TEACHERS ACROSS AMERICA.

TRACK YOUR IMPACT:
WWW.WALGREENS.COM/METOWE

How are we doing?
Enter our monthly sweepstakes for
\$3,000 cash

Visit
WWW.WALGREENSLISTENS.COM

or call toll free
1-800-219-7451
within 72 hours to take a short
survey about this Walgreens visit

SURVEY#
0538-2411-935

PASSWORD
2201-1060-326

For contest rules, see store or
WWW.WALGREENSLISTENS.COM

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT JEFFREY BODIN


DOCTOR K. MCLAUGHLIN, MD

DRUG DESCRIPTION

BIRTH DATE 05/22/97

MEDICATION D-AMPHETAMINE SALT COMBO 30MG TABS

PATIENT ALLERGIES < ALLERGY HISTORY NOT KNOWN >



QUANTITY 90

DIRECTIONS TAKE 1 TABLET BY MOUTH THREE TIMES DAILY

PEACH
FRONT: b / 974
BACK: 30

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. **TELL YOUR DOCTOR:** If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in

life. A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. Erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable. Headache. Trouble sleeping. Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 628188

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2349453-05382 DATE: 11/06/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD
MFG: TEVA
DSB/CMB/CMB/CMB/A.L.AERTKER

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 203114461410168999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 704711777
(985)272-8989

RX # 2349453-05382 DATE: 11/06/20

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

K. MCLAUGHLIN, MD
MFG: TEVA
DSB/CMB/CMB/CMB/A.L.AERTKER

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 203114461410168999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

WAITING

FRI 12:35PM

New


D-AMPHETAMINE SALT COMBO 30MG TABS

00555-0974-02

SAFE

QTY 90

20 DRAM



PEACH
FRONT: b / 974
BACK: 30

DSB/CMB/CMB/CMB/A.L.AERTKER

Med Guide