PATIENT

about side effects. at 1-800-FDA-1088.

advice a

medical effects

r for n

doctor report s

your

Call

JEFFREY BODIN

DOCTOR J. CRUZ, MD

DRUG DESCRIPTION

BIRTH DATE 05/22/97

MEDICATION DYMISTA 137/50 MCG NASAL SPR 120SPR

QUANTITY 23

DIRECTIONS SHAKE LIQUID AND USE 1 SPRAY IN EACH NOSTRIL TWICE DAILY FOR 7 DAYS

PATIENT

INGREDIENT NAME: Azelastine and Fluticasone (a ZEL as teen &

COMMON USES: It is used to ease allergy signs.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to azelastine, fluticasone, or any other part of this drug. TELL any other CCTOR: If you are allergic to any drugs like this one, any other part of this drug. TELL any other CCTOR: If you are allergic to any drugs like this one, any other drugs, so, or other substances. Tell your doctor about the allergy and signs you had, like rash; hives; tiching; shortness of breath; where signs with swelling of face, lips, tongue, or throat; or any other signs. This swelling of face, lips, tongue, or throat; or any other signs. This swelling of face, lips, tongue, or throat; or any other signs. Tell your doctor and interact with other drugs or health problems. Tell your doctor and interact about all of your drugs (prescription or OTC, natural products witamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Do not take this drug by mouth. Use in your nose only. Keep out of your mouth and eyes (may burn). Shake well before use. Prime pump before first use by spraying it 6 times or until you see a fine mist. If you have not used the spray for more than 14 days, you will need to prime the pump with 1 spray for more than 14 days, you will need to prime the pump with 1 spray or until you see a fine mist. Blow your nose before use. Close 1 the strike of the pump with 1 spray or until you see a fine mist. Blow your nose before use. Close 1 the strike of the pump with 1 spray or until you see a fine mist. Blow your nose before use. Close 1 to release spray; by the property of the wall your mouth. Spray up the nose only. Do not spray onto the wall your mouth. Spray up the nose only. Do not spray onto the wall away with two nostriks. If you get this drug in the eyes, fush right away with two nostriks. If you get medical help. HOW DO I STORE AND/OR THROW GUT HIS PRUG? Store upright at room temperature. Do not freeze. Throw away any part not used after 120 sprays. Protect from light. Keep all drugs out of the reach of children and pets. Check with your pharmacist about how to throw out unused drugs. With the missed doses as soon as skip the missed doses as soon as skip the missed doses.

CAUTIONS: Tell all of your health care provider that you take.

skip the missed dose and go back to your normal time. Do not use 2 doses or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. When changing from an oral steroid to another form of a steroid, there may be very bad and sometimes deadly side effects. Signs like weakness, feeling tired, dizziness, upset stomach, throwing up, not thinking clearly, or low blood sugar may happen. Call your doctor right away from the control of the cont

To taik about any risks to your bapy.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; fever whe same; tightness in the chest or throat; trouble breathing or talking, usually loarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of infection like fever, chills, very bad

KFEP OUT OF REACH OF CHILD BEEN ST.

sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Very bad dizziness or passing out. Very upset stomach or throwing up. Feeling very tired or weak. Bad nose irritation. Crusting in the nose. Runny nose. Whistling sound when you breathe. Change in evesight, eye pain, or very bad eye irritation. Bad nosebleeds. Redness or white patches in mouth or throat. Feeling sleepy. WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side medical help if anor fine side effects, your doctor or get medical help if anor fine side effects so the property of the side effects so the property occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1038. You may also report side effects to this PDA at 1-800-FDA-1038. You may also report side effects this this work of the property can be effects at this work of the property side effects.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your durings (prescription, natural products vitamins, OTC) with your drugs with this list to your doctor. Talk with the doctor before starting any this list to your doctor. Talk with the doctor before starting any thing list of all your doctor. Talk with the doctor before starting any companies. One drugs may have another patient information leaflet. Check with your pharmacist; if you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN ille, LA 704711777

RX # 2014427-05382

QTY: 23 New-E

表記 (例)

FRI

New-E

DYMISTA 137/50 MCG NASAL SPR 120SPR 2 REFILLS BEFORE 02/02/19 NDC:00037-0245-23

\$ 0.00

J. CRUZ, MD MFG:MEDA XXX/JJK/JJK/ /B. GUILLORY PLAN: ABHPA GROUP# RX8834 CLAIM REF# 180334153208156999



PH: (985)674-2551

JEFFREY BODIN

rille. I A 704711777

RX # 2014427-05382

DATE: 02/02/18

DYMISTA 137/50 MCG NASAL SPR 120SPR 2 REFILLS BEFORE 02/02/19 NDC:00037-0245-23 OTY: 23

\$ 0.00

J. CRUZ, MD MFG:MEDA XXX/JJK/JJK/ /B. GUILLORY PLAN: ABHPA GROUP# RX8834 CLAIM REF# 180334153208156999

Walgreens

PH: (985)674-2551

Pharmacy use only

1:02PM

DYMISTA 137/50 MCG NASAL SPR 120SPR

00037-**0245**-23

ALPHA

medications or pour not flush unused Do

el società

724.00

88