

AFTER VISIT SUMMARY

Jeffrey Bodin MRN: 1002548110 8/21/2019 9:00 AM University Medical Center ENT Clinic 504-702-3440

ENT Main

Instructions from Laura Pelaez, MD



Today's medication changes

START taking: ofloxacin (OCUFLOX)

Accurate as of August 21, 2019 10:58 AM. Review your updated medication list below.



Pick up these medications from any pharmacy with your printed prescription

ofloxacin



Referrals placed today

Ambulatory Referral to Internal Medicine Where: University Medical Center Family Medicine Clinic Address: 2000 Canal St New Orleans LA 70112-3018 Phone: 504-702-5700 Expires: 8/20/2020 (requested)

Ambulatory Referral to Neurology Where: University Medical Center Neurology Clinic Address: 2000 Canal St New Orleans LA 70112-3018 Phone: 504-962-6210 Expires: 8/20/2020 (requested)

Today's Visit

You saw Laura Pelaez, MD on Wednesday August 21, 2019. The following issues were addressed: Encounter to establish care, Impacted cerumen of right ear, Intractable migraine with aura without status migrainosus, and Primary narcolepsy without cataplexy.



Blood Pressure 119/82



BMI 20.60



Weight 109 lb



Height 5' 1"



Temperature (Oral) 97.9 °F



Pulse 103



Respiration 15



Oxygen Saturation 100%

What's Next

SEP 4 2019 EEG AWAKE AND DROWSY Wednesday September 4 8:30 AM (Arrive by 7:45 AM) University Medical Center Neuro Diagnostics 2000 Canal St New Orleans LA 70112-3018 504-702-5700

JAN 9 2020 Hem Onc Established Patient with Pinki K Prasad, MD Thursday January 9 10:30 AM (Arrive by 10:15 AM) Children's Hospital Hematology and Oncology 200 Henry Clay Ave. New Orleans LA 70118-5720 504-896-9740

FEB 19 2020 Follow Up Appointment with Laura Pelaez, MD Wednesday February 19 8:30 AM (Arrive by 7:45 AM) University Medical Center ENT Clinic 2000 Canal St New Orleans LA 70112-3018 504-702-3440

LCMC Health Patient Portal

Patient is already MyChart active.

ENT - Dr Laura Pelaez

8/21/19

## Need to Schedule an Appointment?

To schedule an appointment, call 504-702-3440. - ENT Main

## ✦ Current Health Issues

Migraine-cluster headache syndrome

Melanoma in situ of left lower leg

Peripheral neuropathy

Inflammatory neuropathy

## Your Allergies

Date Reviewed: 8/21/2019

Allergen

Reactions

Lactose


Nausea And Vomiting










Diarrhea





Latex


Rash

Your Medication List as of August 21, 2019 10:58 AM

 Always use your most recent med list.

	Morning	Around Noon	Afternoon	Evening	Bedtime	As Needed
 <b>AFLURIA QUAD 2018-2019 (PF) 60 mcg</b> (15 mcg x 4)/0.5 mL Syrg injection Generic drug: flu vacc qs2018-19 6mos up(PF) ADM 0.5ML IM UTD						
 <b>ALEVE 220 MG tablet</b> Generic drug: naproxen sodium Take 1,000 mg by mouth 2 (two) times daily with meals						
 <b>azelastine 137 mcg (0.1 %) nasal spray</b> Commonly known as: ASTELIN 1 spray by Nasal route 2 (two) times daily						
 <b>buPROPion 300 MG 24 hr tablet</b> Commonly known as: WELLBUTRIN XL Take 300 mg by mouth daily						
 <b>* dextroamphetamine-amphetamine 20 mg Tab per tablet</b> Commonly known as: ADDERALL Take 30 mg by mouth 3 (three) times daily						
 <b>* dextroamphetamine-amphetamine 30 mg Tab per tablet</b> Commonly known as: ADDERALL TK ONE T PO TID FOR 30 DAYS						
 <b>DYMISTA 137-50 mcg/spray Spry</b> Generic drug: azelastine-fluticasone 1 spray by Nasal route daily						
 <b>fexofenadine 180 MG tablet</b> Commonly known as: ALLEGRA Take 180 mg by mouth daily						
 <b>fluticasone propionate 50 mcg/actuation nasal spray</b> Commonly known as: FLONASE USE ONE SPRAY IEN ONCE D						

		Around				As	
		Morning	Noon	Afternoon	Evening	Bedtime	Needed
 <b>montelukast</b> 10 mg tablet Commonly known as: SINGULAIR Take 10 mg by mouth daily CONTINUE							
 <b>neomycin-polymyxin-hydrocortisone</b> 3.5-10,000-1 mg/mL-unit/mL-% otic suspension Commonly known as: CORTISPORIN Place 3 drops into both ears 5 (five) times daily CONTINUE							
 <b>ofloxacin</b> 0.3 % ophthalmic solution Commonly known as: OCUFLOX Place 4 drops in ear(s) 2 (two) times daily for 10 days Place 4 drops to the right ear twice a day for 10 days Started by: Laura Pelaez, MD START							
 <b>olopatadine</b> 0.2 % Drop ophthalmic solution 1 drop CONTINUE							

 \* This list has 2 medication(s) that are the same as other medications prescribed for you. Read the directions carefully, and ask your doctor or other care provider to review them with you.

# Ear Drops (R Ear) Prescription Info Package

8/21/19

• For more details, visit [Patient Education](#)

Ofloxacin (Ophthalmic) (oh FLOKS a sin)

Brand Names: US Ocuflax

Brand Names: Canada Apo-Ofloxacin; Ocuflax; PMS-Ofloxacin; Sandoz-Ofloxacin

What is this drug used for?

- It is used to treat eye infections.

What do I need to tell my doctor BEFORE I take this drug?

- If you have an allergy to ofloxacin or any other part of this drug.
- If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If you are breast-feeding or plan to breast-feed.

This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Use care when driving or doing other tasks that call for clear eyesight.
- Bright lights may bother you. Wear sunglasses.
- Do not use longer than you have been told. A second infection may happen.
- Very bad and sometimes deadly allergic reactions have rarely happened with other forms of this drug as well as drugs like this one. Talk with the doctor.
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

What are some side effects that I need to call my doctor about right away?

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Change in eyesight, eye pain, or very bad eye irritation.
- Joint pain or swelling.
- A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Burning.
- Eye irritation.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- To gain the most benefit, do not miss doses.
- Use as you have been told, even if your signs get better.
- Use this drug at the same time of day.
- For the eye only.
- Wash your hands before and after use.
- Do not touch the container tip to the eye, lid, or other skin.
- When using, tilt your head back.
- Gently pull down the lower lid and squeeze in how much the doctor told you to use.
- After use, keep your eyes closed. Put pressure on the inside corner of the eye. Do this for 1 to 2 minutes. This keeps the drug in your eye.
- Avoid wearing contacts unless told to wear them by your doctor.

What do I do if I miss a dose?

- Use a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not use 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature. Do not freeze.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor.
- Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

Ear Drops (R Ear) Presc Info Package  
• For wax dis lodging/removal

Page 3 of 3  
8/21/19

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2017-06-15

Copyright

© 2018 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

University Medical Center ENT Clinic  
2000 Canal St  
New Orleans LA 70112-3018  
Phone: 504-702-3440

Date: Aug 21,  
2019


Name: Jeffrey Bodin  
528 BEAU CHENE DR  
MANDEVILLE LA 70471  
985-272-8989

DOB: 05/22/1997

**ofloxacin (OCUFLOX) 0.3 % ophthalmic solution**  
Place 4 drops in ear(s) 2 (two) times daily for 10 days 4 drops to the right  
ear twice a day for 10 days  
End Date: Aug 31, 2019

Disp \*\*\*5 (Five) mL\*\*\* Refill \*\*\*0 (Zero)\*\*\*  
Dispense as Written: No

Associated diagnosis: H61.21

Signature:   
Stephanie Diana Warrington, MD

NPI:1780046631

**VOID**

**VOID**

**VOID**