

2020-02-28 Dr. Linden Appointment Requests

Patient: Jeffrey Thomas Bodin

DOB: 1997-05-22

Request

1. Medication Request (Allergy Medication)

- Medication:

Dymista 13750 MCG Nasal SPR 120SPR

- As of last attempted filling, spring 2018~ approx.
 - Insurance doesn't cover above combination medication.
 - Note: Dymista was covered previously.
 - By my insurance (late 2017 to early 2018).
 - And for years beforehand with my parent's insurance (unknown to late 2017).

1. If Dymista is possible option.

- Then,
 - i. I request a script for above medication.

2. If not possible please substitute for component medications.

- Then,
 - i. I request script for component medications.
 - ii. Or, equal generic substitute (for Dymista).

- Note:

- Following are the component medications:

azelastine hydrochloride

+

fluticasone propionate

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT JEFFREY BODIN	DOCTOR J. CRUZ, MD	DRUG DESCRIPTION
BIRTH DATE 05/22/97		
MEDICATION DYMISTA 137/50 MCG NASAL SPR 120SPR	PATIENT ALLERGIES	
QUANTITY 23		
DIRECTIONS SHAKE LIQUID AND USE 1 SPRAY IN EACH NOSTRIL TWICE DAILY FOR 7 DAYS		

INGREDIENT NAME: Azelastine and Fluticasone (a ZEL as teen & floo TIK a sone)

COMMON USES: It is used to ease allergy signs.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to azelastine, fluticasone, or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. This drug may interact with other drugs or health problems. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Do not take this drug by mouth. Use in your nose only. Keep out of your mouth and eyes (may burn). Shake well before use. Prime pump before first use by spraying it 6 times or until you see a fine mist. If you have not used the spray for more than 14 days, you will need to prime the pump with 1 spray or until you see a fine mist. Blow your nose before use. Close 1 nostril. Tilt your head forward a little. Put nose spray tube into other nostril. While breathing in through the nose, press down once to release spray. Breathe out from your mouth. Spray up the nose only. Do not spray onto the wall joining the two nostrils. If you get this drug in the eyes, flush right away with cool water and get medical help. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store upright at room temperature. Do not freeze. Throw away any part not used after 120 sprays. Protect from light. Keep all drugs in a safe place. Keep up all drugs out of the reach of children and pets. Check with your pharmacist about how to throw out unused drugs.

WHAT DO I DO IF I MISS A DOSE? Use a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not use 2 doses or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. When changing from an oral steroid to another form of a steroid, there may be very bad and sometimes deadly side effects. Signs like weakness, feeling tired, dizziness, upset stomach, throwing up, not thinking clearly, or low blood sugar may happen. Call your doctor right away if you have any of these signs. If you have a bad injury, have surgery, or any type of infection, you may need extra doses of oral steroids. These extra steroids will help your body deal with these stresses. Carry a warning card saying that there may be times when you may need extra steroids. Talk with your doctor if you come into contact with anyone who has chickenpox or measles and you have not had chickenpox, measles, or the vaccines for them. If you have had any recent nose surgery, injury, ulcers, or sores, talk with your doctor. Talk with your doctor before getting any vaccines. Use with this drug may either raise the chance of an infection or make the vaccine not work as well. Talk with your doctor before you drink alcohol or use other drugs and natural products that slow your actions. This drug may raise the chance of cataracts or glaucoma. Talk with the doctor. Have an eye exam as you have been told by your doctor. This drug may cause harm if swallowed. If this drug is swallowed, call a doctor or poison control center right away. This drug may affect growth in children and teens in some cases. They need regular growth checks. Talk with the doctor. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of infection like fever, chills, very bad

sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Very bad dizziness or passing out. Very upset stomach or throwing up. Feeling very tired or weak. Bad nose irritation. Crusting in the nose. Runny nose. Whistling sound when you breathe. Change in eyesight, eye pain, or very bad eye irritation. Bad nosebleeds. Redness or white patches in mouth or throat. Feeling sleepy. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Headache. Nosebleed. Nose irritation. Change in taste. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777

(985)820-4713

RX # 2014427-05382

DATE: 02/02/18

DYMISTA 137/50 MCG NASAL SPR 120SPR

QTY: 23 2 REFILLS BEFORE 02/02/19

New-E NDC:00037-0245-23

\$ 0.00

J. CRUZ, MD
MFG: MEDA
XXXXXXXXXX / B. GULLORY

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 180334153208156999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777

(985)820-4713

RX # 2014427-05382

DATE: 02/02/18

DYMISTA 137/50 MCG NASAL SPR 120SPR

QTY: 23 2 REFILLS BEFORE 02/02/19

New-E NDC:00037-0245-23

\$ 0.00

J. CRUZ, MD
MFG: MEDA
XXXXXXXXXX / B. GULLORY

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 180334153208156999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

FRI 1:02PM
New-E

DYMISTA 137/50 MCG NASAL SPR 120SPR
00037-0245-23
ALPHA

QTY 23

XXXXXXXXXX / B. GULLORY

WIC# 828188 12T

Request

2. Back Pain

- Concerns about pain that is constant when sleeping/laying. Moving etc.
 - Sleeping:
 - Lying in bed facing up.
 - Lying in bed facing down.
 - Lying in bed on either R or L sides.
 - Lying in bed with proper support on entire body.
 - Ie no differing/incorrect elevation of neck/head.
 - Exercising:
 - Exercising and stretching seen no actual change.
 - Pain is still presenting in same manner as previously specified.
 - Have been stretching regularly. (mostly daily to varying degrees).
 - Have purchased exercise equipment. For lower back stretching.
 - No difference thus far as of 2020-02-27.
 - Had since 2020-02-20~ approx.
 - Chair Replacement Repair
 - Have started as of 2020-02-20~ approx ordering items. To supplement the chair's inability to support me properly in current state.
 - Without knowing the actual issue from a medical standpoint. This has proven to be impossible. Please advise on solutions. A diagnosis, even on a cursory level would be greatly helpful.
- In lower back-above waist area.
 - Consistent through all bowel movements. Over course of past 2 months etc. Without any discernable variant with pre/post urination or defecation etc.
- Pain presented itself 6 months ago+ with much less (ie non-constant) degree of frequency.
 - Changed how sat in chair and problem was mitigated.
 - Until now past 3 months. 2019-12-01 to 2020-02-27 approx.

- Chair has degraded to extent that it can no longer be sat in “properly”.
- This is what I believe to be the cause of the current constant back pain.

Request

3. Ophthalmology Appointment/Referral

- I have a current/future appointment set for 2020-03-09 with UMC ophthalmology.
 - With Dr. Peter Robert Kastl. (UMC)
 - 2020-03-09 8:00 am
- I want to ensure this is an Ophthalmology appointment for the melanoma/cancer screening.
 - That I am supposed to be scheduled to receive every 6 months at minimum.
- As I suspect this could be a normal “vision clinic” appointment. For getting new glasses/established with a doctor for this.
 - I am unable to determine if this is the case.
 - Please use the following as guide for correct course of action:
 1. If scheduled appointment is for regular vision purposes.
 - Then,
 - I request a referral for ophthalmology. For melanoma screening.
 2. If scheduled appointment is not for vision. And, is for the cancer.
 - Then,
 - I request a referral for regular vision purposes.

Name: Jeffrey Bodin | DOB: 5/22/1997 | MRN: 1002548110 | PCP: Callie Anne Linden, MD

Appointment Details

New Patient Appointment with Peter Robert Kastl, MD

Monday March 09, 2020

Arrive by 7:15 AM CST

Starts at 8:00 AM CST (30 minutes)

[Add to Calendar](#)

University Medical Center Vision Clinic
2000 Canal St
New Orleans LA 70112-3018
504-702-3950

Prepare for Your Visit

All questionnaires for this appointment will be available for you to answer on Thursday March 05, 2020.

Save time at the clinic by filling out the following questionnaire:

Ophthalmology History (Not Started)

Directions for University Medical Center Vision Clinic

UMCNO Vision Center, 3rd floor of the Clinic (ACB)

Please report to University Medical Center New Orleans. The main patient entrances are on S. Galvez Street and Tulane Avenue. Free patient parking is available in our parking garage at 2001 Tulane Avenue. On the day of your appointment, report to the UMCNO Vision Center on the third floor of the Clinic (ACB). Swipe your valid driver's license or state-issued ID card at the kiosk on your clinic floor. When called, check in with the Patient Access Registrar.

Visit Instructions

Please bring copies of: ---- Medical records ---- Results for labs or tests pertinent to the diagnosis you are being seen for ---- A list of all your medications.

[Cancel Appointment](#)