Walgreens

#05382 4330 HIGHWAY 22 MANDEVILLE, LA 70471 985-674-2551

0392 0041 01/07/2021 11:47 AM 209

FSA RX 2370921 0.00FSA RX 2367983 19.99

SUBTOTAL 19.99 SALES TAX B=4.75% 0.95

TOTAL 20.94 VISA ACCT 1006 20.94

EXPRESS PAY CHANGE .00

TOTAL FSA ITEMS 0.00 TOTAL RX ITEMS

20.94 TOTAL FSA AND RX ITEMS 20.94


APPROVED FSA/HRA AMOUNT 0.00

THANK YOU FOR SHOPPING AT WALGREENS

GET MORE WITH MYWALGREENS, REDEEM WALGREENS CASH REWARDS FOR SOMETHING

EXTRA IN A FUTURE PURCHASE.
RESTRICTIONS APPLY. FOR FULL DETAILS
SEE MYWALGREENS.COM.

RFN# 0538-2410-3921-2101-0703



\$0.29 W CASH REWARDS AVAILABLE

myWalgreens ACCT # ********0353

OPENING BALANCE \$0.10 EARNED THIS VISIT \$0.19

CLOSING BALANCE \$0.29

SEE THE IMPACT OUR BACK TO SCHOOL ME TO WE PROUD SUPPORTERS ARE MAKING ON TEACHERS ACROSS AMERICA.

> TRACK YOUR IMPACT: WWW.WALGREENS.COM/METOWE

How are we doing? Enter our monthly sweepstakes for \$3,000 cash

Visit WWW.WALGREENSLISTENS.COM

> or call toll free 1-800-219-7451

within 72 hours to take a short survey about this Walgreens visit

> SURVEY# 0538-2410-392

> 1210-1070-326

PASSWORD

For contest rules, see store or WWW.WALGREENSLISTENS.COM

DOCTOR K. MCLAUGHLIN, MD

DRUG DESCRIPTION

BIRTH DATE 05/22/97

MEDICATION D-AMPHETAMINE SALT COMBO 30MG TABS

TIMES DAILY

your doctor for medical advice about side effects. may report side effects to FDA at 1-800-FDA-1088.

PATIENT DIRECTIONS TAKE ONE TABLET BY MOUTH THREE

ALLERGIES

< ALLERGY HISTORY NOT KNOWN >

PEACH

FRONT: b / 974

BACK: 30

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. TELL YOUR DOCTOR: If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. TELL YOUR DOCTOR: If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. TELL YOUR DOCTOR: If you have ever had any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid. TELL YOUR DOCTOR: If you have ever had any of these health problems: Drug abuse or stroke. TELL YOUR DOCTOR: If you have kidney disease. TELL YOUR DOCTOR: If you are taking any of these drugs: Acetazolamide or sodium bicarbonate. TELL YOUR DOCTOR: If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen. TELL YOUR DOCTOR: If you are taking any of these drugs: Linezolid or methylene blue. TELL YOUR DOCTOR: If you are breast-feeding. Do not breast-feed while you take this drug. Tills is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN?
Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. WHAT DO I DO IF I MISS A DOSE? Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in

life. A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT INEED TO CALL MY DOCTOR ABOUT RIGHT AWAY?
WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. Erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartheat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects or any other side effects bother you or do not go away. Feeling nervous and excitable. Headache. Trouble sleeping. Constipat

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN

deville, LA 704711777

RX # 2370921-05382

DATE: 01/07/21

QTY:90

D-AMPHETAMINE SALT COMBO 30MG TABS NDC: 00555-0974-02

0.00

K. MCLAUGHLIN, MD MFG:TEVA

PLAN: ABHPA GROUP# RX8834 CLAIM REF# 210074191665104999

Walgreens 4330 HI

PH: (985)674-2551

JEFFREY BODIN

LA 704711777 RX # 2370921-05382

DATE: 01/07/21

D-AMPHETAMINE SALT COMBO 30MG TABS QTY: 90

K. MCLAUGHLIN, MD MFG:TEVA

NO REFILLS NDC:00555-0974-02

> 0.00 PLAN: ABHPA GROUP# RX8834 CLAIM REF# 210074191665104999

Walgreem 4330 HIGHWAY 22 MANDEVILLE, LA PH: (985)674-2551

OUT HERE 🔀

GUT HERE

Pharmacy use only

WAITING

New

THU 11:50AM

D-AMPHETAMINE SALT COMBO 30MG TABS

00555-**0974**-02

SAFE

QTY 90 20 DRAM



FRONT: b / 974

BACK: 30

Med Guide

DSB/BNP/BNP/BNP/B.N.POSTON

flush unused medications or pour down a sink