

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 4330 Highway 22
 Mandeville, LA 70471
 (985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	J. CRUZ, MD	DRUG DESCRIPTION	
BIRTH DATE	05/22/97				
MEDICATION	MONTELUKAST 10MG TABLETS	PATIENT ALLERGIES			
QUANTITY	30				
DIRECTIONS	TAKE 1 TABLET BY MOUTH EVERY DAY				



BEIGE
 FRONT: I
 BACK: 114

INGREDIENT NAME: Montelukast Tablets (MON te LOO kast)

COMMON USES: It is used to ease allergy signs. It is used to prevent exercise-induced breathing problems. It is used to treat or prevent asthma. It may be given to you for other reasons. Talk with the doctor. For asthma: Do not use this drug to treat an asthma attack. Use a rescue inhaler. Talk with your doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to montelukast or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. This drug may interact with other drugs or health problems. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take even during sign-free periods. If working out or playing sports causes signs, use at least 2 hours before doing it. If this drug is for asthma, take in the evening. To gain the most benefit, do not miss doses. Take this drug at the same time of day. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Check with your pharmacist about how to throw out unused drugs. **WHAT DO I DO IF I MISS A DOSE?** Skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Call your doctor right away if your normal dose does not work as well or if you need to use your inhaler more often. This drug is not helpful during an asthma attack. Do not use this drug to stop exercise-induced breathing problems if you are already using it for asthma or allergy. If you have asthma and taking aspirin makes it worse, keep avoiding aspirin and NSAIDs while you take this drug. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side

effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up. Signs of low mood (depression), thoughts of killing yourself, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. Hallucinations (seeing or hearing things that are not there). Change in how you act. Memory problems or loss. Feeling confused. Not able to focus. Bad dreams. Not able to sleep. Sleepwalking. Shakiness. Seizures. Trouble breathing that is new or worse. Fever. Flu-like signs. Sinus pain. Chest pain. A heartbeat that does not feel normal. Any unexplained bruising or bleeding. A burning, numbness, or tingling feeling that is not normal. Ear pain. A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Cough. Loose stools (diarrhea). Headache. Sore throat. Belly pain. Stuffy nose. Runny nose. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE ARE!

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 625188 121

JEFFREY BODIN
 528 Beau Cheve Drive, Mandeville, LA 704711777
 (985)674-4713
RX # 2014426-05382 DATE: 02/02/18

MONTELUKAST 10MG TABLETS
 QTY: 30 2+ REFILLS BEFORE 02/02/19
 New-E NDC: 31722-0726-30

\$ 0.00

J. CRUZ, MD
 MFG: CAMBER
 TRRLTALTA / B. GULLORY

PLAN: ABHPA
 GROUP# RX8834
 CLAIM REF# 180334152468204998

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551

Customer Receipt

JEFFREY BODIN
 528 Beau Cheve Drive, Mandeville, LA 704711777
 (985)674-4713
RX # 2014426-05382 DATE: 02/02/18

MONTELUKAST 10MG TABLETS
 QTY: 30 2+ REFILLS BEFORE 02/02/19
 New-E NDC: 31722-0726-30

\$ 0.00

J. CRUZ, MD
 MFG: CAMBER
 TRRLTALTA / B. GULLORY

PLAN: ABHPA
 GROUP# RX8834
 CLAIM REF# 180334152468204998

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

FRI 1:02PM
 New-E

MONTELUKAST 10MG TABLETS
 31722-0726-30
 FAST RACK

QTY 30
 10 DRAM



BEIGE
 FRONT: I
 BACK: 114

TRRLTALTA / B. GULLORY

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 4330 Highway 22
 Mandeville, LA 70471
 (985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	J. CRUZ, MD	DRUG DESCRIPTION
BIRTH DATE	05/22/97			
MEDICATION	DYMISTA 137/50 MCG NASAL SPR 120SPR			
QUANTITY	23	PATIENT ALLERGIES		
DIRECTIONS	SHAKE LIQUID AND USE 1 SPRAY IN EACH NOSTRIL TWICE DAILY FOR 7 DAYS			

INGREDIENT NAME: Azelastine and Fluticasone (a ZEL as teen & floo TIK a sone)

COMMON USES: It is used to ease allergy signs.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to azelastine, fluticasone, or any other part of this drug. **TELL YOUR DOCTOR:** Tell your doctor if you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. This drug may interact with other drugs or health products. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your other drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Do not take this drug by mouth. Use in your nose only. Keep out of your mouth and eyes (may burn). Shake well before use. Prime pump before first use by spraying it 6 times or until you see a fine mist. If you have not used the spray for more than 14 days, you will need to prime the pump with 1 spray or until you see a fine mist. Blow your nose before use. Close 1 nostril. Tilt your head forward a little. Put nose spray tube into other nostril. While breathing in through the nose, press down once to release spray. Breathe out from your mouth. Spray up the nose only. Do not spray onto the wall joining the two nostrils. If you get this drug in the eyes, flush right away with cool water and get medical help. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store upright at room temperature. Do not freeze. Throw away any part not used after 120 sprays. Protect from light. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Check with your pharmacist about how to throw out unused drugs. **WHAT DO I DO IF I MISS A DOSE?** Use a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not use 2 doses or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. When changing from an oral steroid to another form of a steroid, there may be very bad and sometimes deadly side effects. Signs like weakness, feeling tired, dizziness, upset stomach, throwing up, not thinking clearly, or low blood sugar may happen. Call your doctor right away if you have any of these signs. If you have a bad injury, have surgery, or any type of infection, you may need extra doses of oral steroids. These extra steroids will help your body deal with these stresses. Carry a warning card saying that there may be times when you may need extra steroids. Talk with your doctor if you come into contact with anyone who has chickenpox or measles and you have not had chickenpox, measles, or the vaccines for them. If you have had any recent nose surgery, injury, ulcers, or sores, talk with your doctor. Talk with your doctor before getting any vaccines. Use with this drug may either raise the chance of an infection or make the vaccine not work as well. Talk with your doctor before you drink alcohol or use other drugs and natural products that slow your actions. This drug may raise the chance of cataracts or glaucoma. Talk with the doctor. Have an eye exam as you have been told by your doctor. This drug may cause harm if swallowed. If this drug is swallowed, call a doctor or poison control center right away. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of infection like fever, chills, very bad

sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Very bad dizziness or passing out. Very upset stomach or throwing up. Feeling very tired or weak. Bad nose irritation. Crusting in the nose. Runny nose. Whistling sound when you breathe. Change in eyesight, eye pain, or very bad eye irritation. Bad nosebleeds. Redness or white patches in mouth or throat. Feeling sleepy. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Headache. Nosebleed. Nose irritation. Change in taste. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 828188 12T

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN
 528 Beau Chene Drive, Mandeville, LA 704711777
 (985)820-4713
RX # 2014427-05382 DATE: 02/02/18

DYMISTA 137/50 MCG NASAL SPR 120SPR
 QTY: 23 2 REFILLS BEFORE 02/02/19
 New-E NDC:00037-0245-23

J. CRUZ, MD PLAN: ABHPA
 MFG: MEDA GROUP# RX8834
 XXXJJKJJK / B. GULLORY CLAIM REF# 180334153208156999

Walgreens 4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551 **Customer Receipt**

Pharmacy use only

JEFFREY BODIN
 528 Beau Chene Drive, Mandeville, LA 704711777
 (985)820-4713
RX # 2014427-05382 DATE: 02/02/18

DYMISTA 137/50 MCG NASAL SPR 120SPR
 QTY: 23 2 REFILLS BEFORE 02/02/19
 New-E NDC:00037-0245-23

J. CRUZ, MD PLAN: ABHPA
 MFG: MEDA GROUP# RX8834
 XXXJJKJJK / B. GULLORY CLAIM REF# 180334153208156999

Walgreens 4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551 **Duplicate Receipt**


FRI 1:02PM DYMISTA 137/50 MCG NASAL SPR 120SPR QTY 23
 New-E 00037-0245-23 ALPHA

XXXJJKJJK / B. GULLORY

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 4330 Highway 22
 Mandeville, LA 70471
 (985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT JEFFREY BODIN	DOCTOR A. WEISS, DO	DRUG DESCRIPTION
BIRTH DATE 05/22/97		
MEDICATION D-AMPHETAMINE SALT COMBO 10MG TAB	PATIENT ALLERGIES	
QUANTITY 60		
DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY MORNING AND 1 TABLET AT 2 PM		
		BLUE
		FRONT: b / 972
		BACK: 10

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (am FET a meen/DEX troe am FET a meen)

WARNING: This drug may be habit-forming; avoid long-term use. Use this drug as you were told by your doctor. Tell your doctor if you have a history of drug or alcohol abuse. Misuse of this drug may cause unsafe heart-related side effects or even sudden death. Tell your doctor if you have any heart disease.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with your doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to dextroamphetamine, amphetamine, or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. **TELL YOUR DOCTOR:** If you or a family member has ever had any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma; nervous, anxious, or tense state; or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking acetazolamide. **TELL YOUR DOCTOR:** If you are taking sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. To gain the most benefit, do not miss doses. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, ibuprofen or like products, and some natural products or aids. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place, unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may affect your judgment and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. This drug may be habit-forming with long-term use. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may have some heart tests before starting this drug. Talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Do not switch between this product and other products that have the

same drugs in them without checking with the doctor. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like low mood (depression) or bipolar illness, or if a family member has killed themselves. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like low mood (depression), thoughts of killing yourself, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A very bad and sometimes deadly health problem called serotonin syndrome may happen if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating; hot; very bad diarrhea, upset stomach, or throwing up; or very bad headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING (CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. A fast heartbeat. Very nervous and excitable. Not able to get or keep an erection. Change in sex interest. A heartbeat that does not feel normal. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Dark urine. Not able to pass urine or change in how much urine is passed. Muscle pain or weakness. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with very bad heart problems or heart defects. Call your doctor right away if you have a change in strength on 1 side that is greater than the other, trouble speaking or thinking, eyesight, chest pain or pressure, shortness of breath, or very bad dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Belly pain. Feeling nervous and excitable. Headache. Not hungry. Not able to sleep. Hard stools (constipation). Loose stools (diarrhea). Dizziness. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.**

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 828188 12T

JEFFREY BODIN
 528 Beau Chene Drive, Mandeville, LA 704711777
 (985)620-4713
RX # 2018613-05382 DATE: 02/15/18

D-AMPHETAMINE SALT COMBO 10MG TAB
 QTY: 60 NO REFILLS
 New NDC: 00555-0972-02 \$ 0.00

A. WEISS, DO PLAN: ABHPA
 MFG: TEVA GROUP# RX8834
 NDFMS/MSL/IL RHODES CLAIM REF# 180464462915036999

Walgreens 4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551 Customer Receipt

Pharmacy use only

JEFFREY BODIN
 528 Beau Chene Drive, Mandeville, LA 704711777
 (985)620-4713
RX # 2018613-05382 DATE: 02/15/18

D-AMPHETAMINE SALT COMBO 10MG TAB
 QTY: 60 NO REFILLS
 New NDC: 00555-0972-02 \$ 0.00

A. WEISS, DO PLAN: ABHPA
 MFG: TEVA GROUP# RX8834
 NDFMS/MSL/IL RHODES CLAIM REF# 180464462915036999

Walgreens 4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551 Duplicate Receipt

WAITING D-AMPHETAMINE SALT COMBO 10MG TAB
 THU 12:37PM 00555-0972-02 QTY 60
 New SAFE 10 DRAM

BLUE
 FRONT: b / 972
 BACK: 10

NDFMS/MSL/IL RHODES

Med Guide

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
 4330 Highway 22
 Mandeville, LA 70471
 (985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	A. WEISS, DO	DRUG DESCRIPTION	
BIRTH DATE	05/22/97				
MEDICATION	BUPROPION XL 150MG TABLETS (24 H)	PATIENT ALLERGIES			
QUANTITY	30				
DIRECTIONS	TAKE 1 TABLET BY MOUTH EVERY MORNING				



WHITE
FRONT: WPI 3331

INGREDIENT NAME: Bupropion Extended-Release Tablets (bue PROE pee on)

WARNING: Drugs like this one have raised the chance of suicidal thoughts or actions in children and young adults. The risk may be greater in people who have had these thoughts or actions in the past. All people who take this drug need to be watched closely. Call the doctor right away if signs like low mood (depression), nervousness, restlessness, grouchy/irritability, panic attacks, or changes in mood or actions are new or worse. Call the doctor right away if any thoughts or actions of suicide occur.

COMMON USES: It is used to treat low mood (depression). It is used to treat seasonal affective disorder (SAD). It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to bupropion or any other part of this drug. TELL YOUR DOCTOR: If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. TELL YOUR DOCTOR: If you have ever had seizures. TELL YOUR DOCTOR: If you drink a lot of alcohol and you stop drinking all of a sudden. TELL YOUR DOCTOR: If you use certain other drugs like drugs for seizures or anxiety and you stop using them all of a sudden. TELL YOUR DOCTOR: If you have ever had an eating problem like anorexia or bulimia. TELL YOUR DOCTOR: If you have any of these health problems: Kidney disease or liver disease. TELL YOUR DOCTOR: If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. TELL YOUR DOCTOR: If you are taking any of these drugs: Linezolid or methylene blue. TELL YOUR DOCTOR: If you are taking another drug that has the same drug in it. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Do not take this drug more often than you are told. This may raise the risk of seizures. Be sure you know how far apart to take your doses. Take in the morning if taking once a day. Take with or without food. If you are not able to sleep, do not take this drug too close to bedtime. Talk with your doctor. Swallow whole. Do not chew, break, or crush. To gain the most benefit, do not miss doses. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. If you have trouble swallowing, talk with your doctor. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store in a room in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT TO DO IF I MISS A DOSE?** Skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or activities that call for you to be alert or have clear eyesight until you see how this drug affects you. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not stop taking this drug all of a sudden without calling your doctor. You may have a greater risk of side effects. If you need to stop this drug, you will want to slowly stop it as ordered by your doctor. High blood pressure has happened with this drug. Have your blood pressure checked as you have been told by your doctor. This drug may raise the chance of seizures. The chance may be higher in people who have certain health problems, use certain other drugs, or drink a lot of alcohol. Talk to your doctor to see if you have a greater chance of seizures while taking this drug. Avoid drinking alcohol while taking this drug. Talk with your doctor before you use other drugs and natural products that slow your actions. It may take several weeks to see the full effects. Some people may have a higher chance of eye problems with this drug. Your doctor may want you to have an eye exam to see if you have a higher chance of these eye problems. Call your doctor right away if you have eye pain, change in eyesight, or swelling or redness in or around the eye. If you are taking digoxin, talk with your doctor. You may need to have your blood work checked more closely while you are taking it with this drug. This drug is not approved for use in children. Talk with the doctor. If you are 65 or older, use this drug with care. You could have more side effects. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby. If you smoke: Not all products are approved for use to help stop smoking. Talk with the doctor to make sure that you have the right product. New or worse mental, mood, or behavior problems have happened when bupropion has been used to stop smoking. These problems include thoughts of suicide or killing someone else, depression, suicidal thoughts, anger, anxiety, and anger. These problems have happened in people with and without a history of mental or mood problems. Talk with the doctor.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Change in how you act. Feeling confused. Hallucinations (seeing or hearing things that are not there). If seizures are new or worse after starting this drug. A big weight gain or loss. Chest pain or pressure. Shortness of breath. Ringing in ears. Passing urine more often. Swollen gland. Trouble moving around. Very bad muscle or joint pain. A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have any of these signs: red, swollen, blistered, or peeling skin with or without fever; red or irritated eyes; or sores in your mouth, throat, nose, or eyes. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Dizziness. Headache. Belly pain. Shakiness. Feeling nervous and excitable. Strangely or odd dreams. Upset stomach or throwing up. Hard stools (constipation). Gas. Dry mouth. Not able to sleep. Muscle or joint pain. Nausea or throat irritation. Sweating a lot. Not hungry. A change in weight without trying. For some brands, this is normal and not a cause for concern. For these brands, this is normal and not a cause for concern. If you have questions, talk with your doctor. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain. WIC# 828168 12T

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN
 528 Beau Chene Drive, Mandeville, LA 704711777
 (985)620-4713
RX # 2018612-05382 DATE: 02/15/18

BUPROPION XL 150MG TABLETS (24 H)
 QTY: 30 NO REFILLS - DR. AUTH REQUIRED
 New NDC: 00591-3331-19 \$ 0.00

A. WEISS, DO
 MFG: ACTAVIS
 SMN/RDS/IDE/ /L. RHODES

PLAN: ABHPA
 GROUP# RX8834
 CLAIM REF# 180464461462069999

Walgreens Customer Receipt
 4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551

Pharmacy use only

JEFFREY BODIN
 528 Beau Chene Drive, Mandeville, LA 704711777
 (985)620-4713
RX # 2018612-05382 DATE: 02/15/18

BUPROPION XL 150MG TABLETS (24 H)
 QTY: 30 NO REFILLS - DR. AUTH REQUIRED
 New NDC: 00591-3331-19 \$ 0.00

A. WEISS, DO
 MFG: ACTAVIS
 SMN/RDS/IDE/ /L. RHODES

PLAN: ABHPA
 GROUP# RX8834
 CLAIM REF# 180464461462069999

Walgreens Duplicate Receipt
 4330 HIGHWAY 22 MANDEVILLE, LA 70471
 PH: (985)674-2551

WAITING
 THU 12:37PM
 New

BUPROPION XL 150MG TABLETS (24 H)
00591-3331-19
 ALPHA

QTY 30
10 DRAM

WHITE
FRONT: WPI 3331

SMN/RDS/IDE/ /L. RHODES Med Guide