


YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	C. SWEIDAN, DEN	DRUG DESCRIPTION
BIRTH DATE	05/22/97			 WHITE FRONT: IP33 BACK: 3
MEDICATION	ACETAMINOPHEN/COD #3 (300/30MG) TAB	PATIENT ALLERGIES	<ALLERGY HISTORY NOT KNOWN>	
QUANTITY	12			
DIRECTIONS	TAKE ONE TABLET BY MOUTH EVERY 4 HOURS AS NEEDED FOR PAIN			

INGREDIENT NAME: Acetaminophen and Codeine Tablets (a SEET a MIN oh fen/KOE deen)

WARNING: This drug is a strong pain drug that can put you at risk for addiction, abuse, and misuse. Misuse or abuse of this drug can lead to overdose and death. Talk with your doctor. This drug may cause very bad and sometimes deadly breathing problems. Call your doctor right away if you have slow, shallow, or trouble breathing. The chance of very bad and sometimes deadly breathing problems may be greater when you start this drug, or anytime your dose is raised. Even one especially in children. If this drug is taken by someone else or by accident, medical help right away. This drug has acetaminophen in it. Very bad and sometimes deadly liver problems like the need for a liver transplant have happened with more than 4,000 milligrams of acetaminophen in a day. Also, people who had liver problems were often using more than 1 drug that had acetaminophen in it. Talk with your doctor. This drug has an opioid drug in it. The use of opioid drugs along with has led to very bad side effects. Side effects that have happened include slowed or trouble breathing and deaths. Benzodiazepine drugs include drugs like alprazolam, diazepam, and lorazepam. Benzodiazepine drugs are used to treat many health problems like anxiety, trouble sleeping, or seizures. Talk with the doctor. Many drugs interact with this drug and can raise the chance of side effects like deadly breathing problems. Talk with your doctor and pharmacist to make sure it is safe to use this drug with all of your drugs. Do not take with alcohol or other things that have alcohol. Unsafe and sometimes deadly effects may happen. Caregivers or others need to get medical help right away if the patient does not respond, does not answer or react like normal, or will not wake up. Using this drug for a long time during pregnancy may lead to withdrawal in the newborn baby. This can be life-threatening. Talk with the doctor. Do not give to a child younger than 12 years of age. Children between 12 and 18 years of age who are very overweight or have certain other health problems like sleep apnea or other lung or breathing problems must not use this drug. If your child has been given this drug, ask the doctor for information about the benefits and risks. Some children have had very bad and sometimes deadly breathing problems when using codeine to help them to remove tonsils or adenoids. Do not give to a child younger than 18 years of age who has had surgery to remove tonsils or adenoids. Talk with your child's doctor.

COMMON USES: It is used to ease pain.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to acetaminophen, codeine, or any other part of this drug. TELL YOUR DOCTOR: If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. TELL YOUR DOCTOR: If you have any of these health problems: Lung or breathing problems like asthma, trouble breathing, block or narrowing. TELL YOUR DOCTOR: If you have any of these drugs: Buprenorphine, butorphanol, nalbuphine, or pentazocine. TELL YOUR DOCTOR: If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of these drugs can cause very bad high blood pressure. TELL YOUR DOCTOR: If you are taking any of these drugs: Linezolid or methylene blue. TELL YOUR DOCTOR: If you have been told by your doctor that you are a rapid metabolizer of some drugs. TELL YOUR DOCTOR: If you are breast-feeding or have health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any of your drugs without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take with food if it causes an upset stomach. Drink lots of noncaffeine liquids unless told to drink less liquid by your doctor. Do not take this drug with other strong pain drugs or if you are using a pain patch without talking to your doctor first. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. Do not take more than ordered. If your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug on a regular basis and you stop it all of a sudden, you may have signs of withdrawal. Do not stop taking this drug all of a sudden without calling your doctor. If you have any bad effects, you will be watched closely to make sure you do not become addicted to this drug. HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. WHAT DO I DO IF I MISS A DOSE? If you take this drug on a regular basis, take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses. Many times this drug is taken on an as needed basis. Do not take more often than told by the doctor.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs. This drug may be habit-forming with long-term use. You will be watched closely to make sure you do not misuse, abuse, or become addicted to this drug. Allergic reactions have happened with this drug. Rarely, some reactions can be very bad or life-threatening. Talk with the doctor. Avoid other sources of acetaminophen. Check labels closely. Too much acetaminophen may cause problems. Call your doctor right away if you take more than 4,000 mg (milligrams) of acetaminophen in a day, even if you feel well. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. If you are taking warfarin, talk with your doctor. You may need to have your blood work checked more

closely while you are taking it with this drug. If you are allergic to sulfites, talk with your doctor. Some products have sulfites. Long-term use of an opioid drug like this drug may lead to lower sex hormone levels. This may lead to signs like change in sex ability in men, no menstrual period in women, lowered interest in sex, or fertility problems. Call your doctor if you have any of these signs. This drug may raise the chance of seizures in some people, including people who have had seizures in the past. Talk to your doctor to see if you have a greater chance of seizures while taking this drug. Taking an opioid drug like this drug may lead to a rare but very bad adrenal gland problem. Call your doctor right away if you have very bad dizziness or passing out, very bad upset stomach or throwing up, or if you feel less hungry, very tired, or very weak. If you are 65 or older, use this drug with care. You could have more side effects. The chance of very bad side effects may be higher in children. This may be more likely to happen in children who have breathing problems. Deadly breathing problems have happened with the use of codeine in some children. Talk with the doctor. This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Very bad dizziness or passing out, or pressure or a fast heartbeat. Feeling confused. Very bad constipation. Feeling very tired or weak. Not able to pass urine or change in how much urine is passed. Feeling fever or chills. Sore throat. Mood changes. Very bad headache. Noisy breathing. Seizures. A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; health problem called serotonin syndrome may happen if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea; upset stomach, or throwing up; or very bad headache. WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Constipation. Dizziness. Feeling sleepy. Belly pain. Upset stomach or throwing up. Sweating a lot. Feeling tired or weak. Headache. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188 12T

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN
528 Beau Chene Drive, Mandeville, LA 704711777
(985)272-8989

RX # 2129748-05382

DATE: 01/25/19

ACETAMINOPHEN/COD #3 (300/30MG) TAB
QTY: 12 NO REFILLS - DR. AUTH REQUIRED
New NDC:65162-0033-11

\$ 0.50

C. SWEIDAN, DEN
MFG:AMNEAL
DSB/BJG/BJG/BJG. GULLORY

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 190253120400106999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN
528 Beau Chene Drive, Mandeville, LA 704711777
(985)272-8989

RX # 2129748-05382

DATE: 01/25/19

ACETAMINOPHEN/COD #3 (300/30MG) TAB
QTY: 12 NO REFILLS - DR. AUTH REQUIRED
New NDC:65162-0033-11

\$ 0.50

C. SWEIDAN, DEN
MFG:AMNEAL
DSB/BJG/BJG/BJG. GULLORY

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 190253120400106999

Walgreens


4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

WAITING
FRI 8:54AM
New
ACETAMINOPHEN/COD #3 (300/30MG) TAB
65162-0033-11
CELL 35

QTY 12
10 DRAM



WHITE
FRONT: IP33
BACK: 3

DSB/BJG/BJG/BJG. GULLORY

Med Guide