

Mandeville Pharmacy  
4655 Highway 190  
Mandeville, LA 70448-3582  
Phone: (985) 778-2306  
Fax: (985) 778-2404

Aug 5 2020 9:57AM  
Receipt # 38347  
Cashier: POORNG C  
Drawer #: POS01 - 3275

Rx 43691-00 - JEFFREY 72.99 F  
BODIN

Subtotal 72.99  
Tax Total 3.57

# Items: 1 Total 76.56

Visa Tendered 76.56  
Acct #...1006  
Approval #...025710

F = FLEX/FSA/HRA Eligible  
FSA Eligible \$76.56

Signature required for the following:

- Payment(s) require a signature
- Receipt of Prescriptions
- Receipt of Privacy Practices

**Impacted By  
COVID-19**

Signature Captured Electronically

I agree to pay above total amount according to  
card issuer agreement.

MID: 9139901  
Entry: TAP



Scan This QR Code for Patient Specific  
Medication Information

"Where the Customer is King/Queen"




N0043691-00-00  
JEFFREY BODIN  
RXLOC  
Price 72.99  
08/05/2020  
Sig: \_\_\_\_\_

MA

N0043691 08/05/2020  
JEFFREY BODIN  
Rej: \_\_\_\_\_ N/A: \_\_\_\_\_

MALCOLM ARMSTRONG  
DEXTROAMP-AMPHETAMIN 10 MG TA

 **Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448  
**985-778-2306**  
JEFFREY BODIN  
528 BEAU CHENE DR (985) 272-8989  
MANDEVILLE, LA 70471-1777 PP: RXLO Dr. Kevin McLaughlin  
Rx: N0043691 00 08/05/2020 **\$72.99** Rph: MA  
DEXTROAMP-AMPHETAMIN 10 MG TAB 68382-0952-01 Qty 90



**Mandeville Pharmacy** 1655 Hwy 190 • Mandeville, LA 70448  
**985-778-2306**

**RECEIPT**



Rx: N0043691 00  
JEFFREY BODIN  
528 BEAU CHENE DR

08/05/2020 (985) 272-8989

DOB: 05/22/1997

MANDEVILLE, LA 70471-1777

Dr. Kevin McLaughlin

DEXTROAMP-AMPHETAMIN 10 MG TAB

Qty 90

Auth:

PP: **RXLOC**

NO REFILLS

Rph: MALCOLM ARMSTRONG  
(985) 867-5494  
68382-0952-01  
PHCY ID: 1548616998

Copay

**\$72.99**

**MANDEVILLE PHARMACY**

1655 HIGHWAY 190  
MANDEVILLE, LA 70448-3582

(985) 778-2306

08/05/2020

Rx: N0043691 00  
JEFFREY BODIN

528 BEAU CHENE DR  
MANDEVILLE, LA 70471-1777

(985) 272-8989

Dr. Kevin McLaughlin  
DEXTROAMP-AMPHETAMIN 10 MG TAB

Auth:  
NO REFILLS

Rph: MALCOLM ARMSTRONG  
(985) 867-5494 Qty 90  
68382-0952-01  
PHCY ID: 1548616998

**DUPLICATE  
RECEIPT**

Copay **\$72.99**

PP: **RXLOC**

Call your Doctor for medical advice about side effects.  
You may report side effects to the FDA at: 1-800-FDA-1088.

**We Appreciate Your Business!**

**"Where the Customer is King/Queen"**

**PRESCRIPTION NOTES**

Mandeville Pharmacy  
1655 Highway 190  
Mandeville, LA 70448-3582

For: JEFFREY BODIN  
Rx: 43691  
Date: August 05, 2020  
Message ID: 81842656-407

## **MEDICATION GUIDE**

### **Adderall® (ADD-ur-all) CII**

### **(Dextroamphetamine Saccharate, Amphetamine Aspartate, Dextroamphetamine Sulfate and Amphetamine Sulfate Tablets)**

#### **Rx only**

Read the Medication Guide that comes with Adderall® before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about you or your child's treatment with Adderall®.

#### **What is the most important information I should know about Adderall®?**

**The following have been reported with use of Adderall® and other stimulant medicines.**

#### **1. Heart-Related problems:**

- **sudden death in patients who have heart problems or heart defects**
- **stroke and heart attack in adults**
- **increased blood pressure and heart rate**

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting Adderall®.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with Adderall®.

**Call your doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Adderall®.**

#### **2. Mental (Psychiatric) problems:**

##### **All Patients**

- **new or worse behavior and thought problems**
- **new or worse bipolar illness**
- **new or worse aggressive behavior or hostility**

##### **Children and Teenagers**

- **new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms**

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

**Call your doctor right away if you or your child have any new or worsening mental symptoms or problems while taking Adderall®, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.**

#### **3. Circulation problems in fingers and toes** [Peripheral vasculopathy, including Raynaud's phenomenon]:

- **fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red**

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes.

**Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking Adderall®.**

#### **What is Adderall®?**

Adderall® is a central nervous system stimulant prescription medicine. **It is used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD).** Adderall® may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD. Adderall® should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Adderall® is also used in the treatment of a sleep disorder called narcolepsy.

**Adderall® is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep Adderall® in a safe place to prevent misuse and abuse. Selling or giving away Adderall® may harm others, and is against the law.**

Tell your doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

#### **Who should not take Adderall®?**

**Adderall® should not be taken if you or your child:**

- have heart disease or hardening of the arteries
  - have moderate to severe high blood pressure
  - have hyperthyroidism
  - have an eye problem called glaucoma
  - are very anxious, tense, or agitated
  - have a history of drug abuse
  - are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOI.
  - are sensitive to, allergic to, or had a reaction to other stimulant medicines
- Adderall® is not recommended for use in children less than 3 years old.

**Adderall® may not be right for you or your child. Before starting Adderall® tell your or your child's doctor about all health conditions (or a family history of) including:**

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- liver or kidney problems

- thyroid problems
- seizures or have had an abnormal brain wave test (EEG)
- circulation problems in fingers or toes

Tell your doctor if you or your child are pregnant, planning to become pregnant, or breastfeeding.

### Can Adderall® be taken with other medicines?

**Tell your doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements.** Adderall® and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking Adderall®.

Your doctor will decide whether Adderall® can be taken with other medicines.

### **Especially tell your doctor if you or your child take:**

- anti-depression medicines including MAOIs
- blood pressure medicines
- seizure medicines
- blood thinner medicines
- cold or allergy medicines that contain decongestants
- stomach acid medicines

Know the medicines that you or your child take. Keep a list of your medicines with you to show your doctor and pharmacist.

**Do not start any new medicine while taking Adderall® without talking to your doctor first.**

### How should Adderall® be taken?

- **Take Adderall® exactly as prescribed.** Your doctor may adjust the dose until it is right for you or your child.
- Adderall® tablets are usually taken two to three times a day. The first dose is usually taken when you first wake in the morning. One or two more doses may be taken during the day, 4 to 6 hours apart.
- Adderall® can be taken with or without food.
- From time to time, your doctor may stop Adderall® treatment for a while to check ADHD symptoms.
- Your doctor may do regular checks of the blood, heart, and blood pressure while taking Adderall®. Children should have their height and weight checked often while taking Adderall®. Adderall® treatment may be stopped if a problem is found during these check-ups.
- **If you or your child take too much Adderall® or overdoses, call your doctor or poison control center right away, or get emergency treatment.**

### What are possible side effects of Adderall®?

See **“What is the most important information I should know about Adderall®?”** for information on reported heart and mental problems.

### **Other serious side effects include:**

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision

### **Common side effects include:**

- stomach ache
- decreased appetite
- nervousness

Adderall® may affect your or your child’s ability to drive or do other dangerous activities.

Talk to your doctor if you or your child have side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your doctor or pharmacist

for more information.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

### How should I store Adderall®?

- Store Adderall® in a safe place at room temperature, 20° to 25°C (68° to 77°F).
- **Keep Adderall® and all medicines out of the reach of children.**

### General information about Adderall®

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Adderall® for a condition for which it was not prescribed. Do not give Adderall® to other people, even if they have the same condition. It may harm them and it is against the law. This Medication Guide summarizes the most important information about Adderall®. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Adderall® that was written for healthcare professionals. For more information about Adderall®, please contact Teva Pharmaceuticals at 1-888-838-2872.

### What are the ingredients in Adderall®?

**Active Ingredient:** dextroamphetamine saccharate, amphetamine aspartate monohydrate, dextroamphetamine sulfate, USP, amphetamine sulfate, USP

**Inactive Ingredients:** lactitol, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, and other ingredients.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

[Teva Select Brands, Horsham, PA 19044](#)

Division of Teva Pharmaceuticals USA

Revised October 2015

# Patient Education - Dextroamp-Amphetamin 10 Mg Tab

**CONSUMER INFORMATION USE AND DISCLAIMER** This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Dextroamphetamine and Amphetamine Tablets

PRONUNCIATION (deks troe am FET a meen & am FET a meen)

BRAND NAME(S): Adderall

**WARNING** This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take for longer than you have been told by your doctor. Use only as you were told. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death.

**WHAT IS THIS DRUG USED FOR?** It is used to treat attention deficit problems with hyperactivity.; It is used to treat narcolepsy.; It may be given to you for other reasons. Talk with the doctor.

**WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG?** If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.; If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics.; If you have any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid.; If you have ever had any of these health problems: Drug abuse or stroke.; If you have kidney disease.; If you are taking any of these drugs: Acetazolamide or sodium bicarbonate.; If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen.; If you are taking any of these drugs: Linezolid or methylene blue.; If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

**WHAT ARE SOME THINGS I NEED TO KNOW OR DO WHILE I TAKE THIS DRUG?** Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. ; Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.; If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered.; If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug.; You may need to have some heart tests before starting this drug. If you have questions, talk with your doctor.; This drug may cause high blood pressure.; Check blood pressure and heart rate as the doctor has told you.; Have blood work checked as you have been told by the doctor. Talk with the doctor.; This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.; Do not take antacids with this drug.; Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.; New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life.; A severe and sometimes deadly problem called serotonin syndrome may happen if you take this drug with certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache.; This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.; Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child.; Talk with your doctor before you drink alcohol.; Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

**WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.; Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.; Not able to get or keep an erection.; Change in sex interest.; Seizures.; Trouble controlling body movements.; Restlessness.; Change in eyesight.; For males, erections (hard penis) that happen often or that last a long time.; Change in color of hands or feet from pale to blue or red.; Numbness, pain, tingling, or cold feeling of the hands or feet.; Any sores or wounds on the fingers or toes.; Muscle pain or weakness, dark urine, or trouble passing urine.; Pain when passing urine.; Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out.

**WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable.; Headache.; Trouble sleeping.; Constipation, diarrhea, stomach pain, upset stomach, throwing up, or feeling less hungry.; Feeling dizzy, tired, or weak.; Dry mouth.; Bad taste in your mouth.; Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

**HOW IS THIS DRUG BEST TAKEN?** Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.; Take with or without food.; Take last dose of the day at least 4 hours before bedtime.

**WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it.; If it is close to the time for your next dose, skip the missed dose and go back to your normal time.; Do not take 2 doses at the same time or extra doses.

**HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.; Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.; Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

**GENERAL DRUG FACTS** If your symptoms or health problems do not get better or if they become worse, call your doctor. ; Do not share your drugs with others and do not take anyone else's drugs.; This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If

you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.; If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Copyright 2020 UpToDate, Inc. All Rights Reserved. (Issue 20.3.3.005)