


YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT JEFFREY BODIN	DOCTOR A. WEISS, DO	DRUG DESCRIPTION
BIRTH DATE 05/22/97		
MEDICATION D-AMPHETAMINE SALT COMBO 30MG TABS	PATIENT ALLERGIES <ALLERGY HISTORY NOT KNOWN>	
QUANTITY 90		PEACH
DIRECTIONS TAKE 1 TABLET BY MOUTH THREE TIMES DAILY		FRONT: b / 974
		BACK: 30

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. Use this drug only as you were told by your doctor. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to dextroamphetamine, amphetamine, or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. **TELL YOUR DOCTOR:** If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. **TELL YOUR DOCTOR:** If you have any of these health problems: Glaucoma; nervous, anxious, or tense state; or overactive thyroid. **TELL YOUR DOCTOR:** If you have ever had any of these health problems: Drug abuse or stroke. **TELL YOUR DOCTOR:** If you have kidney disease. **TELL YOUR DOCTOR:** If you are taking acetazolamide. **TELL YOUR DOCTOR:** If you are taking sodium bicarbonate. **TELL YOUR DOCTOR:** If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. **TELL YOUR DOCTOR:** If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. This drug may be habit-forming with long-term use. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may have some heart tests before starting this drug. Talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Talk with the doctor. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, ibuprofen or like products, and some natural products or aids. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you or a family member have any mental or mood problems like low mood (depression) or bipolar illness, or if a family member has killed themselves. Call your doctor right away if you have hallucinations; change in the way you act; or signs of mood changes like low mood

(depression), thoughts of killing yourself, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A very bad and sometimes deadly health problem called serotonin syndrome may happen if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea, upset stomach, or throwing up; or very bad headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Not able to get or keep an erection. Change in sex interest. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Muscle pain or weakness, dark urine, or trouble passing urine. Pain when passing urine. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with some heart problems or heart defects. Call your doctor right away if you have a fast, slow, or abnormal heartbeat; weakness on 1 side of the body; trouble speaking or thinking; change in balance; drooping on 1 side of the face; change in eyesight; chest pain or pressure; shortness of breath; or severe dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Feeling nervous and excitable. Headache. Not hungry. Trouble sleeping. Diarrhea or constipation. Stomach pain. Feeling dizzy, tired, or weak. Dry mouth. Bad taste in your mouth. Weight loss. Upset stomach or throwing up. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188 12T

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777
(985)272-8989

RX # 2205306-05382 DATE: 09/14/19

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

A. WEISS, DO
MFG: TEVA
MOB/JWT/JWT/JWT/J.W.TALLEY

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 192573251592217999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551

Customer Receipt

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777
(985)272-8989

RX # 2205306-05382 DATE: 09/14/19

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90 NO REFILLS
New NDC: 00555-0974-02

\$ 0.00

A. WEISS, DO
MFG: TEVA
MOB/JWT/JWT/JWT/J.W.TALLEY

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 192573251592217999

Walgreens

4330 HIGHWAY 22 MANDEVILLE, LA 70471
PH: (985)674-2551


Duplicate Receipt

Pharmacy use only

SAT 10:28AM
New

D-AMPHETAMINE SALT COMBO 30MG TABS
00555-0974-02
SAFE

QTY 90
20 DRAM



PEACH
FRONT: b / 974
BACK: 30

MOB/JWT/JWT/JWT/J.W.TALLEY

Med Guide