PATIENT

ALLERGIES

MEDICATION D-AMPHETAMINE SALT COMBO 30MG TABS

QUANTITY

DIRECTIONS TAKE 1 TABLET BY MOUTH THREE

TIMES DAILY

DOCTOR A. WEISS, DO

DRUG DESCRIPTION

PEACH

FRONT: b / 974 BACK: 30

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (deks troe am FET a meen & am FET a meen)

WARNING: This drug has a risk of abuse and misuse. This drug may also be habit-forming if taken for a long time. Do not take longer than you have been told by your doctor. Use this drug only as you were told by your doctor. Tell your doctor if you have ever abused or been addicted to any drugs or alcohol. Misuse of this drug may cause heart-related side effects or even sudden death. If you have any questions, talk with your doctor.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

Present USING THIS MEDICINE: WHAT DO INEED TO TELL MY DOCTOR BEFORE USING THIS MEDICINE: WHAT DO INEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG'T ELL YOUR DOCTOR: If YOU have an allergy to dextroamphetamine, amphetamine, or any other part of this drug. Tags. You dextroamphetamine, amphetamine, or any other part of this drug. Tags. You so the substances. Tell your doctor about the allergy and whet signs you had, like rash, hives; itching; shortness of breath; wheezing; cough; swelling of face, ips, tongue, or throat; or any other signs. TELL YOUR DOCTOR: If you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. TELL YOUR DOCTOR: If you have any of these health problems; Drug abuse or stroke. TELL YOUR DOCTOR: If you have sidney disease. FELL YOUR DOCTOR: If you have kidney disease. FELL YOUR DOCTOR: If you have kidney disease. FELL YOUR DOCTOR: If you are taking and the problems of the part of the problems. Brug abuse or stroke. TELL YOUR DOCTOR: If you have kidney disease. FELL YOUR DOCTOR: If you are taking as the problems of these drugs are retaind rugs used for low mood (depression) like isocarboxazid, phenelzine or team creation drugs used for low mood (depression) like isocarboxazid, phenelzine or the last 14 days. Taking used for Parkinson's disease of those drugs can cause very bad high blood pressure. TELL YOUR DOCTOR: If you are breast feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems. You are taking any of these drugs: Linezolid or methylene blue. TELL YOUR DOCTOR: If you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the day at least 4 hours before bedtime. To gain the most benefit, do not miss doses. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. Talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, ibuprofen or like products, and some natural products or aids. HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. WHAT DO I DO IFI MSS. A COSE Take a missed dose as soon as you think about it. If it is close to the total control of the same time or extra doses.

the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. This drug may be habit-forming with long-term use. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may have some heart tests before starting this drug. Talk with your doctor. This drug may avenue to slowly stop this drug. You may have some heart tests before starting this drug. Talk with your doctor. This drug may avenue heart part as the doctor has told you. Talk with the doctor. Have blood pressure and heart rate as the doctor has told you. Talk with the doctor. Have blood work checked as you have been told by the doctor. Talk with the doctor. Have blood works that you take this drug. Do not take antaclds with this drug. Do not switch between this product and other products that have the same drugs in them without of the product and other products that have the same drugs in them vinthus of the product and other products that have the same drugs in them without or a family member have happened with this drug. Do not switch between this product and other products that have the same drugs in them without or a family member have the same drugs in them without or a family member have the same drugs in them without or a family member have the same drugs in the workers that you doctor if you doctor if you act a family member have the same drugs in the way you act; or signs of mood changes like low

benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect. Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or taking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. A fast heartbeat. Very nervous and excitable. Not able to get or keep an erection. Change in sex interest. A heartbeat that does not feel normal. Seizures. Trouble controlling body, movements. Restlessness. Change in eyesight. For males, erections (hard penis)

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Dark urine. Not able to pass urine or change in how much urine is passed, Pain when passing urine. Muscle pain of the passed pain the passing urine. Muscle pain of the passed pain the passing urine. Muscle pain of the passing urine of the passing urine. Muscle pain of the passing urine of the passing urine and the passing urine. Muscle pain or passing the passing urine and the passing urine passing u

< ALLERGY HISTORY NOT KNOWN >

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor, Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

JEFFREY BODIN

RX # 2182729-05382

DATE: 07/05/19

D-AMPHETAMINE SALT COMBO 30MG TABS QTY:90

NO REFILLS

New

NDC:00555-0974-02

\$ 0.00

A. WEISS, DO MFG:TEVA MOJ/KLT/KLT/ANM/E.K.DAVIS PLAN: ABHPA GROUP# RX8834 CLAIM REF# 191865662847083999

Walgreens

PH: (985)674-2551

JEFFREY BODIN

e, LA 704711777

RX # 2182729-05382

DATE: 07/05/19

D-AMPHETAMINE SALT COMBO 30MG TABS

QTY: 90

NO REFILLS NDC:00555-0974-02

A. WEISS, DO MFG:TEVA MOJ/KLT/KLT/ANM/E.K.DAVIS

PLAN: ABHPA GROUP# RX8834 CLAIM REF# 191865662847083999

Walgreens

PH: (985)674-2551

\$ 0.00

Pharmacy use only

3:58PM

WAITING

FRI

New

D-AMPHETAMINE SALT COMBO 30MG TABS

00555-**0974**-02

SAFE

QTY 90 20 DRAM

PEACH

FRONT: b / 974 BACK: 30

Med Guide

flush unused medications or pour