

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location

2880 Highway 190
Mandeville, LA 70471
(985)624-8548

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT JEFFREY BODIN

DOCTOR PBR: R. DEWITZ


DRUG DESCRIPTION

BIRTH DATE 05/22/97

MEDICATION HYDROCODONE/ACETAMINOPHEN 5-325 TB

QUANTITY 28

PATIENT ALLERGIES



DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 6 HOURS AS NEEDED FOR PAIN FOR UP TO 7 DAYS

WHITE
FRONT: 2172

INGREDIENT NAME: Hydrocodone and Acetaminophen Tablets and Capsules (HYE droe KOE done/a SEET a MIN on ten)

WARNING: This drug may be habit-forming with long-term use. This drug is a strong pain drug that can put you at risk for addiction, abuse, and misuse. Misuse or abuse of this drug can lead to overdose and death. Talk with your doctor. You will be watched closely to make sure you do not misuse, abuse, or become addicted to this drug. This drug may cause very bad and sometimes deadly breathing problems. Call your doctor right away if you have slow, shallow, or trouble breathing. The chance of very bad and sometimes deadly breathing problems may be greater when you first start this drug or anytime your dose is raised. Even one dose of this drug may be deadly if it is taken by someone else or by accident, especially in children. If this drug is taken by someone else or by accident, get medical help right away. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. This drug has an opioid drug in it. The use of opioid drugs along with a benzodiazepine drug or other drugs that may make you drowsy or slow your actions has led to very bad side effects. Side effects that have happened include slowed or trouble breathing and deaths. Benzodiazepine drugs include drugs like alprazolam, diazepam, and lorazepam. Benzodiazepine drugs are used to treat many health problems like anxiety, trouble sleeping, or seizures. Talk with the doctor. Many drugs interact with this drug and can raise the chance of side effects like deadly breathing problems. Talk with your doctor and pharmacist to make sure it is safe to use this drug with all of your drugs. Do not take with alcohol or products that have alcohol. Unsafe and sometimes deadly effects may happen. Get medical help right away if you feel very sleepy, very dizzy, or if you pass out. Caregivers or others need to get medical help right away if the patient does not respond, does not answer or react like normal, or will not wake up. This drug has acetaminophen in it. Very bad and sometimes deadly liver problems like the need for a liver transplant have happened with acetaminophen use. Most of the time, liver problems have happened in people taking more than 4,000 milligrams of acetaminophen in a day. Also, people who had liver problems were often using more than 1 drug that had acetaminophen in it. Talk with your doctor. Using this drug, for a long time during pregnancy may lead to withdrawal in the newborn baby. This can be life-threatening. Talk with the doctor.

COMMON USES: It is used to ease pain.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: If you have an allergy to hydrocodone, acetaminophen, or any other part of this drug. **TELL YOUR DOCTOR:** If you are allergic to any drugs, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. **TELL YOUR DOCTOR:** If you have any of these health problems: Lung or breathing problems like asthma, trouble breathing, or sleep apnea; high levels of carbon dioxide in the blood; or stomach or bowel block or narrowing. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Buprenorphine, bupropion, nalbuphine, or pentazocine. **TELL YOUR DOCTOR:** If you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. **TELL YOUR DOCTOR:** If you are taking any of these drugs: Linezolid or methylene blue. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC products, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as directed by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take with food if it causes an upset stomach. **HOW DO I STORE AND/OR THROW OUT THIS DRUG?** Store at room temperature. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep away from the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. **WHAT DO I DO IF I MISS A DOSE?** If you take this drug on a regular basis, take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses. Many times this drug is taken on an as needed basis. Do not take more often than told by the doctor.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you. To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs. Allergic reactions have happened with this drug. Rarely, some reactions can be very bad or life-threatening. Talk with the doctor. Do not take this drug with other strong pain drugs or if you are using a pain patch without talking to your doctor first. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. If you have been taking this drug on a regular basis and you stop it all of a sudden, you may have signs of withdrawal. Do not stop taking this drug all of a sudden without calling your doctor. Tell your doctor if you have any bad effects. Avoid other sources of acetaminophen. Check labels closely. Too much acetaminophen may cause problems. Call your doctor right away if you take more than 4,000 mg (milligrams) of acetaminophen in a day, even if you feel well. Long-term use of an opioid drug like this drug may lead to lower sex hormone levels. This may lead to signs like change in sex ability in men, no menstrual period in women, lowered interest in sex, or fertility problems. Call your doctor if you have any of these signs. Taking an opioid drug like this drug may lead to a rare but very bad adrenal gland problem. Call your doctor right away if you have very bad dizziness or passing out, very bad upset stomach or throwing up, or if you feel less hungry, very tired, or very weak. This drug may raise the chance of seizures in some people, including people who have had seizures in the past. Talk to your doctor to see if you have a greater chance of seizures while taking this drug. If you are 65 or older, use this drug with care. You could have more side effects. Use with care in children. Talk with the doctor. This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away. Tell your doctor if you are breast-feeding or plan to breast-feed. This drug passes

into breast milk and may harm your baby.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Very bad dizziness or passing out. Feeling very sleepy. Trouble breathing, slow breathing, or shallow breathing. Feeling confused. Trouble walking. Very hard stools (constipation). Very bad belly pain. Feeling very tired or weak. Not able to pass urine or change in how much urine is passed. Hearing loss. Change in hearing. Chest pain or pressure or a fast heartbeat. Fever or chills. Sore throat. Mood changes. Very bad headache. Trouble passing urine. Any unexplained bruising or bleeding. Change in eyesight. Seizures. A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes. A very bad and sometimes deadly health problem called serotonin syndrome may happen if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea, upset stomach, or throwing up; or very bad headache. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Hard stools (constipation). Dizziness. Feeling sleepy. Upset stomach or throwing up. Feeling tired or weak. Headache. Belly pain. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 828188 12T

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777
(985)520-4713

RX # 3951878-03440

DATE: 05/28/18

HYDROCODONE/ACETAMINOPHEN 5-325 TB

QTY: 28 NO REFILLS

New NDC: 00591-2172-05

\$ 0.00

PBR: R. DEWITZ
MFG: ACTAVIS
SPB/SPB/SPB /S. BOLLICH

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 181486944304165996

Walgreens

2880 HIGHWAY 190 MANDEVILLE, LA 70471
PH: (985)624-8548

Customer Receipt

JEFFREY BODIN

528 Beau Chene Drive, Mandeville, LA 704711777
(985)520-4713

RX # 3951878-03440

DATE: 05/28/18

HYDROCODONE/ACETAMINOPHEN 5-325 TB

QTY: 28 NO REFILLS

New NDC: 00591-2172-05

\$ 0.00

PBR: R. DEWITZ
MFG: ACTAVIS
SPB/SPB/SPB /S. BOLLICH

PLAN: ABHPA
GROUP# RX8834
CLAIM REF# 181486944304165996

Walgreens

2880 HIGHWAY 190 MANDEVILLE, LA 70471
PH: (985)624-8548


Duplicate Receipt

Pharmacy use only

MON 8:42PM
New

HYDROCODONE/ACETAMINOPHEN 5-325 TB
00591-2172-05
SAFE

QTY 28
10 DRAM



WHITE
FRONT: 2172

SPB/SPB/SPB /S. BOLLICH

Med Guide