


YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	W. TERRAL JR, MD	DRUG DESCRIPTION	
BIRTH DATE	05/22/97				PEACH
MEDICATION	D-AMPHETAMINE SALT COMBO 20MG TABS				FRONT: b / 973
QUANTITY	45				BACK: 20
DIRECTIONS	TAKE 1 TABLET BY MOUTH THREE TIMES DAILY				

INGREDIENT NAME: Dextroamphetamine and Amphetamine Tablets (am FET a meen/DEX troe am FET a meen)

WARNING: This drug may be habit-forming; avoid long-term use. Use this drug as you were told by your doctor. Tell your doctor if you have a history of drug or alcohol abuse. Misuse of this drug may cause unsafe heart-related side effects or even sudden death. Tell your doctor if you have any heart disease.

COMMON USES: It is used to treat attention deficit problems with hyperactivity. It is used to treat narcolepsy. It may be given to you for other reasons. Talk with the doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR: if you have an allergy to dextroamphetamine, amphetamine, or any other part of this drug. TELL YOUR DOCTOR: if you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. TELL YOUR DOCTOR: if you or a family member have any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics. TELL YOUR DOCTOR: if you have any of these health problems: Glaucoma; nervous, anxious, or tense state; or overactive thyroid. TELL YOUR DOCTOR: if you have ever had any of these health problems: Drug abuse or stroke. TELL YOUR DOCTOR: if you have kidney disease. TELL YOUR DOCTOR: if you are taking acetazolamide. TELL YOUR DOCTOR: if you are taking sodium bicarbonate. TELL YOUR DOCTOR: if you have taken certain drugs used for low mood (depression) like isocarboxazid, phenelzine, or tranylcypromine or drugs used for Parkinson's disease like selegiline or rasagiline in the last 14 days. Taking this drug within 14 days of those drugs can cause very bad high blood pressure. TELL YOUR DOCTOR: if you are taking any of these drugs: linezolid or methylene blue. TELL YOUR DOCTOR: if you are breast-feeding. Do not breast-feed while you take this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take last dose of the drug at least 4 hours before bedtime. To gain the most benefit, do not miss doses. Keep taking this drug as you have been told by your doctor or your health care provider, even if you feel well. Talk with your doctor or health care provider, even if you raise blood pressure. These include: cough or cold drugs; diet pills, stimulants, ibuprofen or like products; some natural products or aids. HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Check with your pharmacist about how to throw out unused drugs. WHAT DO I DO IF I MISS A DOSE? Take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Avoid driving and doing other tasks or actions that you for you to be alert until you see how this drug affects you. If you have been taking this drug for a long time or at high doses, it may not work as well and you may need higher doses to get the same effect. This is known as tolerance. Call your doctor if this drug stops working well. Do not take more than ordered. This drug may be habit-forming with long-term use. If you have been taking this drug for many weeks, talk with your doctor before stopping. You may want to slowly stop this drug. You may have some heart tests before starting this drug. Talk with your doctor. This drug may cause high blood pressure. Check blood pressure and heart rate as the doctor has told you. Talk with the doctor. Have blood work checked as you have been told by the doctor. Talk with the doctor. This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug. Do not take antacids with this drug. Do not switch between this product and other products that have the same drugs in them without checking with the doctor. New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell your doctor if you

or a family member have any mental or mood problems like low mood (depression) or bipolar illness, or if a family member has killed themselves. Call your doctor right away if you have hallucinations: change in the way you act; or signs of mood changes like low mood (depression), thoughts of killing yourself, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. A very bad and sometimes deadly health problem called serotonin syndrome may happen if you take this drug with drugs for depression, migraines, or certain other drugs. Call your doctor right away if you have agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; very bad diarrhea, upset stomach, or throwing up; or very bad headache. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child. Talk with your doctor before you drink alcohol. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? **WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness; passing out, or change in eyesight. A fast heartbeat. Very nervous and excited. Change in sex ability. Lowered interest in sex. A heartbeat that does not feel normal. Seizures. Trouble controlling body movements. Restlessness. Change in eyesight. For males, erections (hard penis) that happen often or that last a long time. Change in color of hands or feet from pale to blue or red. Numbness, pain, tingling, or cold feeling of the hands or feet. Any sore or wound on the fingers or toes. Dark urine. Not able to pass urine or change in how much urine is passed. Muscle pain or weakness. Heart attacks, strokes, and sudden deaths have happened in adults taking this drug. Sudden deaths have also happened in children with very bad heart problems or heart defects. Call your doctor right away if you have a change in strength on 1 side that is greater than the other, trouble speaking or thinking, change in balance, drooping on 1 side of the face, change in eyesight, chest pain or pressure, shortness of breath, or very bad dizziness or passing out. **WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG?** All drugs may cause side effects. However, many people who get medical help if any of these side effects or any other side effects bother you or do not go away: Belly pain. Feeling nervous and excitable. Headache. Not hungry. Not able to sleep. Hard stools (constipation). Loose stools (diarrhea). Dizziness. Dry mouth. Bad taste in your mouth. Weight loss. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at http://www.fda.gov/medwatch.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with your doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. This drug comes with an extra rating fact sheet called a Medication Guide. Read it with care. Read it each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.

JEFFREY BODIN
528 Beau Chene Dr, Mandeville, LA 7047111777
(985)264-1080

RX # 1967915-05382 DATE: 09/19/17

D-AMPHETAMINE SALT COMBO 20MG TABS
QTY: 45 NO REFILLS
New NDC: 00555-0973-02

W. TERRAL JR, MD PLAN: PERX
MFG: TEVA GROUP# BSLA
KJWIFZAFZAABC.B. GULLORY CLAIM REF# FFMWKWH



\$ 15.00


Customer Receipt

Walgreens
4330 HIGHWAY 22, MANDEVILLE, LA 70471
PH: (985)674-2551

Pharmacy use only

WAITING
TUE 2:21PM
New

D-AMPHETAMINE SALT COMBO 20MG TABS
00555-0973-02
SAFE



PEACH
FRONT: b / 973
BACK: 20

KJWIFZAFZAABC.B. GULLORY

Med Guide

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

JEFFREY BODIN
528 Beau Chene Dr, Mandeville, LA 7047111777
(985)264-1080

RX # 1967915-05382 DATE: 09/19/17

D-AMPHETAMINE SALT COMBO 20MG TABS
QTY: 45 NO REFILLS
New NDC: 00555-0973-02

W. TERRAL JR, MD PLAN: PERX
MFG: TEVA GROUP# BSLA
KJWIFZAFZAABC.B. GULLORY CLAIM REF# FFMWKWH

\$ 15.00

Duplicate Receipt

Walgreens
4330 HIGHWAY 22, MANDEVILLE, LA 70471
PH: (985)674-2551


Call your doctor for medical advice about side effects.
Do not flush unused medications or pour down a sink or drain.

WIC# 216382 12T

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location
4330 Highway 22
Mandeville, LA 70471
(985)674-2551

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	JEFFREY BODIN	DOCTOR	W. TERRAL JR, MD	DRUG DESCRIPTION	
BIRTH DATE	05/22/97				
MEDICATION	MONTELUKAST 10MG TABLETS				
QUANTITY	90	PATIENT ALLERGIES			

DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY DAY

BEIGE
FRONT: I
BACK: 114

INGREDIENT NAME: Montelukast Tablets (MON te LOO kast)

COMMON USES: It is used to ease allergy signs. It is used to prevent exercise-induced breathing problems. It is used to treat or prevent asthma. It may be given to you for other reasons. Talk with the doctor. For asthma: Do not use this drug to treat an asthma attack. Use a rescue inhaler. Talk with your doctor.

BEFORE USING THIS MEDICINE: WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? TELL YOUR DOCTOR if you have an allergy to montelukast or any other part of this drug. TELL YOUR DOCTOR: If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. This drug may interact with other drugs or health problems. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

HOW TO USE THIS MEDICINE: HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Take with or without food. Take even during sign-free periods. If working out or playing sports causes signs, use at least 2 hours before doing it. If this drug is for asthma, take in the evening. To gain the most benefit, do not miss doses. Take this drug at the same time of day. Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well. HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature. Protect from light. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Check with your pharmacist about how to throw out unused drugs. WHAT DO I DO IF I MISS A DOSE? Skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses.

CAUTIONS: Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Call your doctor right away if your normal dose does not work as well or if you need to use your inhaler more often. This drug is not helpful during an asthma attack. Do not use this drug to stop exercise-induced breathing problems if you are already using it for asthma or allergy. If you have asthma and taking aspirin makes it worse, keep avoiding aspirin and NSAIDS while you take this drug. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

POSSIBLE SIDE EFFECTS: WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side

effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up. Signs of low mood (depression), thoughts of killing yourself, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life. Hallucinations (seeing or hearing things that are not there). Change in how you act. Memory problems or loss. Feeling confused. Not able to focus. Bad dreams. Not able to sleep. Sleepwalking. Shakiness. Seizures. Trouble breathing that is new or worse. Fever. Flu-like signs. Sinus pain. Chest pain. A heartbeat that does not feel normal. Any unexplained bruising or bleeding. A burning, numbness, or tingling feeling that is not normal. Ear pain. A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes. WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Cough. Loose stools (diarrhea). Headache. Sore throat. Belly pain. Stuffy nose. Runny nose. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

OVERDOSE: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

ADDITIONAL INFORMATION: If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

Call your doctor for medical advice about side effects
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382 121

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 7047111777
(985)264-1080

RX # 1945691-05382

DATE: 10/01/17

MONTELUKAST 10MG TABLETS

QTY: 90 1 REFILL BEFORE 07/07/18

Refill NDC: 31722-0726-30

\$ 33.01

W. TERRAL JR., MD PLAN: PERX
MFG: CAMBER GROUP# BSLA
ZZZ / / J.W.TALLEY CLAIM REF# R3MNER1

Walgreens

4330 HIGHWAY 22, MANDEVILLE, LA 70471

PH: (985)674-2551

Pharmacy use only

Customer Receipt

JEFFREY BODIN

528 Beau Chene Dr, Mandeville, LA 7047111777
(985)264-1080

RX # 1945691-05382

DATE: 10/01/17

MONTELUKAST 10MG TABLETS

QTY: 90 1 REFILL BEFORE 07/07/18

Refill NDC: 31722-0726-30

\$ 33.01

W. TERRAL JR., MD PLAN: PERX
MFG: CAMBER GROUP# BSLA
ZZZ / / J.W.TALLEY CLAIM REF# R3MNER1

Walgreens

4330 HIGHWAY 22, MANDEVILLE, LA 70471

PH: (985)674-2551

Duplicate Receipt

Pharmacy use only

MON 12:00PM Refill
MONTELUKAST 10MG TABLETS
31722-0726-30
FAST RACK

QTY 90
10 DRAM
BEIGE
FRONT: I
BACK: 114